


Muon combined reconstruction in the ATLAS detector with first data





María Moreno Llácer (IFIC, Valencia)
María J. Costa (IFIC, Valencia)
XXXII Bienal de Física,
Ciudad Real, September 2009

Contents

- Introduction: ATLAS
- Muon reconstruction in the ATLAS detector
- Commissioning of ATLAS
- Analysis:
 - Samples of cosmic data
 - Correlations inner detector (ID) & muon spectrometer (MS)
 - Performance of the combined (ID+MS) reconstruction
 - Muon charge ratio measurement
 - Strategy and systematics
 - Energy loss in the calorimeters
- Conclusions

Introduction:

A Toroidal LHC Apparatus (ATLAS)



Inner Detector (ID), $B=2T$:

- Silicon pixels and strips (SCT)
- Transition radiation tracker TRT (separates e/π)

Calorimeters:

- EM: Pb-LAr accordion shape
- HAD: Fe/scintillator (central), Cu/W-LAr (forward)

Muon Spectrometer (MS), $B=0.5T/\text{air-core}$:

- 4 technologies: MDTs and CSCs (precision), RPCs and TGCs (trigger)
- air-core toroid system

Length ~ 46 m

Diameter ~ 25 m

Weight ~ 7000 tons


Muon reconstruction in ATLAS

LHC physics requires an excellent identification and measurement of muons in a large energy range:

~GeV (B physics) and ~TeV (new physics)

→ It is essential combine information of the sub-detectors in ATLAS

Standalone tracking





Muon spectrometer
Calorimeters
Inner detector

Not enough:

- There are dead regions
- ID and calorimeters info needed
- low p_t muons:
 - signal only in first chambers
 - worst p_t resolution
- combined (MS+ID) reconstruction needed

María Moreno Llácer (IFIC,CSIC-UV)


Combined tracking: ID+MS




Commissioning of ATLAS

Test beams


Installation of the sub-detectors



Cosmic data taking:
Milestone weeks



Sept. 2008:
single beam



Cosmic data taking
- Fall 2008
- June/July 2009




Fall 2009: Single beam
2010: Proton collisions









In fall 2008, several weeks of combined cosmic ray data taking involving the whole ATLAS detector took place. Several million tracks with and without B field were reconstructed. This work will study the performance of the **Global χ^2 combined (ID+MS) tracking** with cosmic data.

Correlations between Inner Detector & Muon Spectrometer tracks




Correlation ID and MS tracks

Tracks are characterized
by 5 parameters: (d_0 , z_0 , ϕ , θ , q/p)



- ID and Muon tracks show the same acceptance since events with ID tracks are selected.
- Good correlation between the parameters measured in both sub-detectors ID and MS are synchronized and aligned \rightarrow combined tracking can be attempted.
- The data/MC agreement is fairly good.

Performance of the COMBINED TRACKING (Inner Detector + Muon Spectrometer)



Performance of the combined tracking


Most of the tracks come through the **barrel** and **cross the whole detector**.

In each half (top or bottom) there are:



Transition Radiation Tracker (TRT):
36 straw layers

Semi-Conductor Tracker (SCT):
4 barrel layers

Silicon pixels:
3 barrel layers




Hits in the pixels




$$\sigma_{\text{residuals}} = \sqrt{\sigma_{\text{detector}}^2 + \sigma_{\text{track}}^2 + \sigma_{\text{alignment/calibration}}^2}$$

SCT residuals



- The data/MC agreement is fairly good for all sub-detectors with the current alignment constants.
- Slightly more associated hits in simulation than in data.

Charge muon ratio measurement



Charge muon ratio measurement



- Cosmic rays are primarily protons
 - Excess of π^+ / K^+ over π^- / K^- in shower development
- Expect to see some excess of positive muons.
- Useful and fun exercise.

Measurement strategy:

tracks measured as positive

tracks measured as negative

- Raw Ratio, Center of ATLAS
 - P'/N' (versus p')
- Corrected Ratio, Center of Detector
 - P/N (versus p') = $\epsilon^- / \epsilon^+ \cdot f(P', N', c)$
- “Unsmeared” Ratio, Center of Detector
 - $p' \rightarrow p$
- Ratio at sea level
 - wrong charge probability




Understanding the systematics

- A physics quality measurement of the charge muon ratio requires to understand:

- trigger acceptance, efficiency & timing
- ID & MS tracking efficiency
- charge confusion
- track parameters resolution
- extrapolate to surface (to compare with PDG)



Charge dependent acceptance




Muons are bend out of ATLAS depending on its charge and position
 → charged bias acceptance.

Charge confusion

OS: opposite sign
 SS: same sign





Energy lost in the calorimeters



Momentum difference ID-MS & energy loss

- When a muon traverses the detector material, it undergoes successive deflections and a loss of energy.
- Cosmic muons lose energy in the calorimeters primarily through ionization.
- This energy deposited in the calorimeters is the difference in the cosmic muon momentum before (at the end of the muon system) and after (at the entrance of the inner detector) crossing them.



- The energy deposited in the calorimeters is smaller than the ID-MS momentum difference.
- Expected difference due to dead material: ~20%
- Present measured discrepancy: ~200 MeV (~7%)
- Next data-reprocessing: expect a discrepancy of ~21%

Conclusiones

- The identification and measurement of muons is fundamental for the physics studies in ATLAS.
- The muon reconstruction algorithms have been verified for cosmic rays data using information of all ATLAS sub-detectors (ID, calorimeters and MS) and the performance of the combined tracking has been studied. The reconstruction has been done using the GRID facilities.
- Muon energy loss studies and a measurement of the μ^+/μ^- asymmetry are in progress.
- Data/MC comparisons have been performed and show relatively good agreement with the current alignment and calibration constants.

Thank you very much! 😊

BACK-UP


Samples of cosmic data analyzed and event selection

Four “golden” runs with similar detector and readout conditions from late October 2008 have been used in the analysis. Most of the sub-detectors were operated at full coverage and the magnetic field was on.

This work will focus on the performance of the combined tracking:

- Track reconstruction in the ID
- Track reconstruction in the MS

→ **Global χ^2 combined tracking**



→ **Event selection & track quality cuts:**

Select events with:

- 1 track in the ID, ≥ 1 in the MS & and 1 combined
- Triggered by the muon system
- Select projective muons (impact parameters $< 30\text{cm}$)
- Track quality cuts (#hits) applied