



Property measurements of the Higgs boson in the $\gamma\gamma$ final state with the ATLAS detector at the LHC

Main SM production modes:




H → $\gamma\gamma$ decays:



Reconstructing H → $\gamma\gamma$ @ ATLAS


- Simple topology:
 - 2 isolated energetic photons
- Good discrimination from hadronic background:
 - Show shapes in EM calorimeter
- Narrow signal peak:
 - Good mass resolution (FWHM ~ 2-5 GeV)



Calibration and PID efficiency of photons

- Calorimeter non-uniformities and layer intercalibration correction
- EM cluster energy correction via MVA regression
- E-scale and resolution extracted with Z → ee

Energy scale uncertainty: 0.03%-0.05% for 40 GeV E_T electron, 0.2-0.5% for photons




Energy resolution uncertainty: 10% better resolution for H → $\gamma\gamma$, 5-10% for 10-45 GeV electron.

3 different methods for PID efficiency measurement


- photons from Z → $\gamma\ell\ell$
- extrapolation from electron in Z → ee decays
- Matrix method based on photon purity.

Time	PID unc.
July 2012	10.8%
Dec. 2012	5.3%
Mar. 2013	2.4%
Now	1.0%



Significant improvement achieved

Mass of the Higgs boson



Channel	Mass measurement [GeV]
H → $\gamma\gamma$	125.98 ± 0.42 (stat) ± 0.28 (syst) = 125.98 ± 0.50
H → ZZ* → 4 ℓ	124.51 ± 0.52 (stat) ± 0.06 (syst) = 124.51 ± 0.52
Combined	125.36 ± 0.37 (stat) ± 0.18 (syst) = 125.36 ± 0.41


Previous results in H → $\gamma\gamma$: 126.8 ± 0.2 (stat) ± 0.7 (syst)

Analysis Strategy

- Definition of the Fiducial region
 - 2 isolated photons with $|\eta| < 2.37$
 - $p_{T\gamma 1}(p_{T\gamma 2})/m_{\gamma\gamma} > 0.35(0.25)$
 - $105 < m_{\gamma\gamma} < 160$ GeV
- Observables
 - Higgs boson kinematic: $P_{T\gamma\gamma}, |\eta_{\gamma\gamma}|, P_{Tt}$
 - Jet activity: $N_{jets}, P_{Tj1}, H_{Tjet}$, etc.
 - VBF-sensitive variables: $\Delta\phi_{\gamma\gamma jj}, m_{jj}, P_{THjj}$, etc.
 - Spin-CP sensitive variables: $\cos\theta^*, \Delta\phi_{jj}, \Delta y_{jj}$

Fiducial and Differential X-sec measurements

Direct comparison with the state-of-art theoretical predictions



Signal extraction (for each observable)

- S+B simultaneous fit to all bins for an observable of interest with the Higgs mass fixed to combination measurement.





illustration of simultaneous fit for N_{jets}



Bin-by-bin Correction Factors



Here, v_i and c_i are the extracted signal yield and correction factor in the i^{th} bin, respectively

Compatibility

- Agreement is quantified:
 - 1st / 2nd moment



Broadly in line with the theoretical expectations

