



Searches for Higgs bosons decaying to lepton pairs with the CMS detector

Jan Steggemann for the CMS Collaboration

CERN

Abstract

Searches for Higgs bosons decaying to lepton pairs are summarised using the run 1 dataset recorded by the CMS detector. The first part of this summary is devoted to the search for Higgs bosons decaying to tau pairs. The search is carried out using all six tau-pair final states, with the tau decaying to a muon, an electron, or hadrons. Evidence for Higgs boson decays to taus is reported, and constraints on the couplings of the Higgs boson to fermions and bosons are derived. Then, searches for Higgs boson decays to muons and electrons are also discussed. The last part discusses a search for lepton-flavour violating Higgs boson decays to a muon and a tau, with the tau being reconstructed in the decay to an electron or hadrons. The analysis places the best limits on the respective Yukawa couplings to date while showing a slight excess of 2.5 standard deviations above the standard model expectation.

Keywords: Standard-model Higgs bosons, taus, lepton flavour violation

1. Introduction

At the time of the previous ICHEP conference in 2012, the discovery of a particle was reported with mass near 125 GeV and properties consistent with those of the standard model (SM) Higgs boson. While the initial and subsequent analyses measured various of the production and decay properties to be consistent with the predictions of the SM [1, 2, 3, 4, 5], measurements in different decay channels and with higher precision are necessary to probe the compatibility with the SM predictions and the mechanism of fermion mass generation. The searches for Higgs boson decays to leptons address several of the remaining open questions, most importantly whether the Higgs boson decays to fermions at all.

The first goal of the search for $H \rightarrow \tau\tau$ decays is to answer this question, i.e. to test whether the Higgs boson decays to taus, leptons, and fermions [6]. In the context of all SM Higgs boson decay modes, the $H \rightarrow \tau\tau$ final state is expected to provide the most stringent constraints on the combined couplings to fermions. The $H \rightarrow \tau\tau$ analysis also provides high sensitivity to vector boson fusion (VBF) production.

While searches for $H \rightarrow \mu\mu$ and $H \rightarrow ee$ decays are

(at least currently) not sensitive to the event rates predicted by the SM, they allow to test whether the Higgs boson couplings to leptons are proportional to its mass, as predicted by the SM, or whether the couplings are flavour-universal, i.e. the same as for taus. Furthermore, $H \rightarrow \mu\mu$ decays probe the second-generation Yukawa couplings, and an enhanced $H \rightarrow \mu\mu$ decay rate may be a sign of physics beyond the SM [7].

Lepton-flavour violating (LFV) decays of the Higgs boson to leptons may occur in several extensions of the SM, e.g. in models with multiple Higgs doublets, composite Higgs models, and Randall-Sundrum models [8]. While the branching fraction to muons and electrons is severely constrained from the measurement of $\mu \rightarrow e$ transitions, $B(H \rightarrow \mu e) < O(10^{-8})$, constraints on the branching fractions $B(H \rightarrow \mu\tau)$ and $B(H \rightarrow e\tau)$ are significantly weaker, $B < O(10\%)$. The first dedicated search for $H \rightarrow \mu\tau$ decays is discussed here and is expected to significantly improve the limit on $B(H \rightarrow \mu\tau)$.

In the following, Higgs boson decay modes are always given explicitly, e.g. $H \rightarrow \tau\tau$ denotes Higgs boson decays to two taus with unspecified tau decays, and $H \rightarrow \tau\mu$ denotes the LFV direct decay of the Higgs

boson to a tau and a muon. Reconstructed lepton final states, on the other hand, are indicated by only the two final-state leptons, e.g. μe for an electron-muon final state. Reconstructed hadronic tau decays are denoted as τ_h . In all considered analyses, the final-state leptons are required to have opposite charge, which is therefore not explicitly noted.

This summary is organised as follows. Section 2 describes the search for $H \rightarrow \tau\tau$ decays in the context of the SM. The following section 3 gives a brief overview of the searches for $H \rightarrow \mu\mu$ and $H \rightarrow ee$ decays. Finally, the search for LFV $H \rightarrow \tau\mu$ decays is reported in section 4, after which an overall summary is given.

2. Evidence for the Higgs boson decaying to a pair of taus

The branching fraction of Higgs boson decays to taus is the fourth-largest in the SM, $B(H \rightarrow \tau\tau) = 6.3\%$ for $m_H = 125$ GeV. The two taus decay further and can be reconstructed in the following 6 di-tau final states in order of decreasing branching fraction, $\tau_h\tau_h, \mu\tau_h, e\tau_h, \mu e, \mu\mu, ee$, which are accompanied by neutrinos. All 6 tau-pair final states are considered.

Events are recorded using trigger requirements based on reconstructed electron, muon, and τ_h candidates. The leptons are required to have low transverse momentum (p_T) from nearby reconstructed photons, neutral, and charged hadrons reconstructed by the particle flow algorithm, as quantified by isolation criteria, both on trigger level and in the baseline event selection. Different p_T thresholds are applied for the different leptons depending on the final state and the corresponding trigger requirements. The missing transverse energy (E_T^{miss}) is reconstructed using a multivariate regression technique, which improves both the separation against backgrounds with high- p_T neutrinos and the di-tau mass reconstruction.

The relevant Higgs boson production modes for this analysis are gluon fusion, VBF, and the associated production with a W or Z boson. The different production modes are targeted by splitting the events into different categories based on the number of additional jets and leptons, and their properties. Gluon fusion production is targeted by requiring no or at least one additional jet (“0-jet/1-jet”). In the latter case, the Higgs boson may obtain a significant amount of p_T , thereby increasing the separation against the relevant background processes. VBF production is addressed by requiring two additional jets compatible with VBF production, i.e. with a large rapidity gap and without a central jet (“VBF tag”). Finally, the associated production with a W or Z boson

is targeted by requiring one or two additional muons or electrons in the final state that are compatible with coming from a W or Z boson decay, respectively.

Additional event selection criteria are applied for a number of final states, including multivariate classifiers to separate against $t\bar{t}$ background in the $e\mu$ channel and against $Z \rightarrow \ell\ell$ background in the ee and $\mu\mu$ channels. To reject W+jets events, a low transverse mass $m_T < 30$ GeV is required in the $\mu\tau_h$ and $e\tau_h$ channels.

Because of the high branching fraction, the high production cross section, and the good separation against background processes, the most important final states are $\tau_h\tau_h, \mu\tau_h, e\tau_h$, and μe , in particular in the 1-jet and VBF-tag categories. In these leptonic final states, the final observable is the reconstructed di-tau mass, calculated from the two reconstructed leptons and the E_T^{miss} vector. Figure 1 shows, on the one hand, the distribution of the reconstructed visible mass from the two reconstructed leptons and, on the other hand, the reconstructed mass using the so-called SVFit algorithm. The SVFit algorithm is based on a likelihood fit that takes the matrix elements for the tau decay as well as the experimental resolutions of the different input variables into account. Using the SVFit algorithm significantly improves the separation against the major background, $Z \rightarrow \tau\tau$, and leads to an increase of the final expected significance by $\sim 40\%$ compared to using only the visible mass reconstruction.

To optimise the sensitivity of the search, the events are further divided into sub-categories to increase the separation of signal and background events using (i) the p_T of either the τ_h in final states with one τ_h , of the μ in the μe final state, or of the lepton with highest p_T in the $\mu\mu$ and ee final states, (ii) the p_T of the reconstructed di-tau system, and (iii) the mass of and the $\Delta\eta$ between the two jets in the VBF-tag categories.

Figure 2 shows the combined mass distribution in the four most important final states. The events are weighted by the signal over signal-plus-background ratios in a central 68% interval of the signal distributions in each event category. The figure also shows the distributions of the major backgrounds. The overall largest background is from $Z \rightarrow \tau\tau$ production. It is modelled using a so-called “embedding” technique where the reconstructed muons in $Z \rightarrow \mu\mu$ data are replaced by generated taus. Among the other important backgrounds are W+jets production, estimated using a control region with high transverse mass, and QCD multijet production, estimated using control regions with same-sign leptons and/or inverted lepton isolation requirements.

The final results of the analysis are obtained from a combined likelihood fit of all event categories, includ-





Figure 1: Top figure: Normalised distribution of the visible invariant mass of the reconstructed $\mu\tau_h$ pair in the $\mu\tau_h$ final state. Bottom figure: According distribution of the reconstructed di-tau mass using the SVFit algorithm.




Figure 2: Combined mass distribution in the four most important final states targeting gluon fusion and VBF production. The events are weighted by the signal over signal-to-background ratio in a central 68% interval of the signal distribution in each category. The background distributions are obtained from the final combined fit.




Figure 3: Observed and expected local p-values and significances in terms of standard deviations as a function of m_H .

149 ing all relevant systematic uncertainties. One important
 150 feature of the fit is it constrains several of the systematic
 151 uncertainties. For example, the tau energy scale and tau
 152 reconstruction efficiencies are constrained by the posi-
 153 tion and height of the Z boson mass peaks in the differ-
 154 ent lepton final states and event categories, in particular
 155 also from the distributions in the 0-jet categories that
 156 have a large contribution from $Z \rightarrow \tau\tau$ events.

157 The observed and expected significance of the anal-
 158 ysis are shown in figure 3. The expected significance
 159 for $m_H = 125$ GeV amounts to 3.7 standard deviations

160 (σ), and it is the highest among all m_H for two reasons.
 161 On the one hand, the mass of 125 GeV provides a bet-
 162 ter separation against the Z boson mass peak than lower
 163 m_H values. On the other hand, the cross section times
 164 branching fraction falls as as function of m_H , giving a
 165 higher expected yield than higher m_H values. The cor-

166 responding observed significance at 125 GeV is 3.2σ ,
 167 constituting evidence for $H \rightarrow \tau\tau$ decays. To be only
 168 sensitive to $H \rightarrow \tau\tau$ decays, Higgs boson decays in the
 169 only other relevant $H \rightarrow WW$ channel are treated as a
 170 background in the calculation of the significances and
 171 best-fit signal strength modifiers in terms of the SM ex-
 172 pectation (μ).




Figure 4: Measured μ values for different final states based on the number of jets and the lepton multiplicity, and combined expected and observed μ value for $m_H = 125$ GeV.

173 The μ values for the different jet and lepton multi-
 174 plicity final states are given in figure 4. The combined
 175 best-fit μ value is 0.78 ± 0.27 . Both the combined and the
 176 individual μ values are compatible with the SM expec-
 177 tation of $\mu = 1$. As can be seen from the uncertainties,
 178 the largest sensitivity comes from the 1-jet and VBF-tag
 179 categories.

180 Figure 5 shows a likelihood scan for the fermion and
 181 vector boson couplings. The parameters κ_f and κ_V de-
 182 note the ratio between the measurements and SM pre-
 183 dictions of the Higgs boson couplings to fermions and
 184 vector bosons, respectively. In this case, the contri-
 185 bution from $H \rightarrow WW$ is treated as a signal process.
 186 The results are compatible with the SM expectation of
 187 $\kappa_f = \kappa_V = 1$.

188 Finally, the results from the $H \rightarrow \tau\tau$ search are com-
 189 bined with a search for $H \rightarrow b\bar{b}$ [9, 10]. The combined
 190 and individual likelihood scans as a function of μ are
 191 shown in figure 6. The observed significance of the
 192 $H \rightarrow b\bar{b}$ search is 2.1σ , and the combined observed sig-
 193 nificance amounts to 3.8σ with an expectation of 4.4σ .
 194 This result constitutes strong evidence for the direct de-




Figure 5: Likelihood scan for the combined couplings to fermions κ_f and vector bosons κ_V .




Figure 6: Likelihood scan as a function of the signal strength modifier μ for the $H \rightarrow \tau\tau$ analysis, the $H \rightarrow b\bar{b}$ analysis, and their combination.

195 cay of the Higgs boson to fermions.

196 3. Search for Higgs boson decays to muon or elec- 197 tron pairs

198 The $H \rightarrow \mu\mu$ branching fraction of 2.2×10^{-4} is
 199 among the smallest that is expected to be eventually
 200 measurable at the LHC. The $H \rightarrow \mu\mu$ search profits from

the high experimental resolution of the reconstructed di-muon mass of a few GeV [11]. However, the SM decay width of the Higgs boson is still small compared to the experimental resolution.

The analysis strategy is similar to the $H \rightarrow \tau\tau$ search. The gluon fusion and VBF production modes are targeted by requiring additional jets in the final state. In addition, the events are split into categories according to the p_T of the di-muon system, the properties of the jets in the 2-jet category, and according to the detector regions in which the two muons are reconstructed. The latter make use of the different experimental resolutions of the reconstructed di-muon mass for muons from different detector regions.

The combined di-muon mass distributions are shown in figure 7 top, weighted for the ratio of the signal and signal-plus-background distributions in the different event categories. The background distribution is estimated using an analytic fit function, and, amongst others, systematic uncertainties are estimated by changing the used fit function.

Figure 7 bottom shows the upper limits on the $H \rightarrow \mu\mu$ cross section times branching fraction in terms of the SM expectation. The combined expected limit is 5.1 times the SM expectation, while the observed 95% confidence level (CL) limit is 7.4 times the SM expectation.

For $H \rightarrow ee$, the inaccessibly small branching fraction expected in the SM is $B = 5 \times 10^{-9}$. The search follows the same strategy as the $H \rightarrow \mu\mu$ search. The resulting 95% CL upper limits on the $H \rightarrow ee$ cross section times branching fraction are shown in figure 8. They are of similar size as the limits on the $H \rightarrow \mu\mu$ cross section times branching fraction.

Contrasted with the evidence for $H \rightarrow \tau\tau$ decays, the absence of a signal in the $H \rightarrow \mu\mu$ and $H \rightarrow ee$ searches shows that the branching fraction of the Higgs boson decay to the different leptons is not identical, i.e. the decay is not universal in lepton flavour.

4. Search for LFV decays of the Higgs boson to a muon and a tau

The previous best constraints on the $H \rightarrow \tau\mu$ branching fraction are from a reinterpretation of an ATLAS $H \rightarrow \tau\tau$ search with only 7 TeV LHC data, which are more stringent than the indirect limits from $\tau \rightarrow \mu\gamma$ decays [12]. This suggests that the first dedicated search discussed here is expected to significantly improve these constraints [8]. The $H \rightarrow \tau\mu$ search considers decays of the tau to an electron or to hadron, leading to the two lepton final states $\mu\tau_h$ and μe ; the $\mu\mu$ final state is




Figure 7: Top figure: Combined distribution of the reconstructed di-muon mass. The events are weighted by the ratio of the signal and signal-plus-background expectations in each individual category. Bottom figure: Upper limits on the ratio of the cross section times $B(H \rightarrow \mu\mu)$ in terms of the SM cross section times branching fraction as a function of m_H .

not considered because of the large background from $Z \rightarrow \mu\mu$ production.

The analysis strategy is similar to the one in the search for SM Higgs boson decays to tau pairs in the corresponding $\mu\tau_h$ and μe final states. The gluon fusion production mode is targeted with two event categories requiring either 0 or 1 jet in the final state in addition to the Higgs boson decay products, and VBF production is targeted with another event category requiring two jets




Figure 8: Upper limits on the $H \rightarrow ee$ cross section times branching fraction as a function of m_H .

259 consistent with coming from VBF production, leading
 260 to 3 event categories per lepton final state, i.e. 6 in total.
 261 Since the lepton final states are identical, the basic
 262 event selection is the same as in the two considered decay
 263 channels.

264 However, a number of kinematic distributions are different
 265 as the $H \rightarrow \tau\mu$ decay only involves neutrinos in the decay
 266 of the tau. This implies that (a) the muon p_T distribution
 267 is harder than in the $H \rightarrow \tau\tau$ case, (b) there is a large
 268 azimuthal angle between the muon and the sum of the neutrinos
 269 that are reconstructed as the E_T^{miss} vector in the x-y plane,
 270 and (c) there is a small azimuthal angle between the visible
 271 tau decay products, i.e. either the electron or the τ_h , and the
 272 invisible ones being reconstructed as the E_T^{miss} vector, since
 273 they originate from the same high- p_T tau.

274 To make use of these differences, additional event selection
 275 criteria are applied in each of the categories based on the
 276 p_T of the leptons, the differences in azimuthal angle between
 277 the final state leptons and between the electron and the
 278 E_T^{miss} vector in the μe final state, and the transverse mass
 279 values calculated from either of the leptons and the E_T^{miss}
 280 vector. The additional selection criteria are optimised separately
 281 in each of the 6 event categories.

282 As the final observable, the so-called “collinear mass” is used.
 283 Since there is only one tau in the event with high p_T compared
 284 to its mass, the approximation is made that the neutrino momenta,
 285 measured as the E_T^{miss} vector, and the momenta of the visible
 286 tau decay products,

289 are in the same direction as the tau momentum.

290 The backgrounds are also estimated with similar methods as in
 291 the SM $H \rightarrow \tau\tau$ search. Due to the tighter event selection,
 292 two important differences arise. First, the $Z \rightarrow \tau\tau$ can be
 293 suppressed more efficiently, meaning that other sources of
 294 background are most relevant, in particular W +jets, multijet,
 295 and $t\bar{t}$ events. Second, due to this suppression, the 0-jet
 296 categories also provide high sensitivity.




Figure 9: Upper limits on the branching fraction of LFV $H \rightarrow \tau\mu$ decays. The limits are shown separately for the 6 different event categories and their combination.

298 Upper limits on the branching fraction of $H \rightarrow \tau\mu$
 299 decays are obtained with the CL_S method using a combined
 300 likelihood fit. The resulting limits are shown in figure 9,
 301 both for the 6 event categories and their combination. The
 302 expected upper limit on the $H \rightarrow \tau\mu$ branching fraction is
 303 0.75%, while the observed limit is 1.57%. With respect to the
 304 background-only expectation, a slight excess is observed. The
 305 background-only p-value is 0.007 for $m_H = 126$ GeV, corresponding
 306 to a significance of 2.5σ . The best-fit branching fraction is
 307 measured to be $B(H \rightarrow \tau\mu) = (0.89^{+0.40}_{-0.37})\%$.

308 Finally, the results are translated into limits on the flavour-
 309 violating Yukawa couplings, $|Y_{\mu\tau}|$ and $|Y_{\tau\mu}|$, as shown in
 310 figure 10. The presented search improves the previous best
 311 limits on the Yukawa couplings by an order of magnitude.




Figure 10: Constraints on the flavour-violating Yukawa coupling $Y_{\mu\tau}$, with the expected limits given by the red line and the observed limits as the full black line. The previous best direct experimental limits from $\tau \rightarrow \mu\gamma$ decays are given by the white line and the middle blue-green area, and the limits from a re-interpretation of an ATLAS $H \rightarrow \tau\tau$ search are drawn as a yellow line.

5. Conclusion

In the search for $H \rightarrow \tau\tau$ decays, evidence for Higgs boson decays to tau pairs is obtained, with an observed significance of 3.2 standard deviations given an expectation of 3.7 standard deviations. The measured event rates, mass, and couplings to fermions and vector bosons are consistent with expectations from the SM. The combination with the search for Higgs boson decays to b quarks yields strong evidence for Higgs boson decays to fermions. Tight constraints are also derived on the $H \rightarrow \mu\mu$ and $H \rightarrow ee$ branching fractions, showing that the Higgs boson couplings to leptons are not flavour-universal.

The search for LFV provides the most stringent limits $H \rightarrow \tau\mu$ decays and the $Y_{\tau\mu}$ Yukawa coupling to date. Results with LHC run 2 data will resolve whether the slight excess that is observed can be attributed to a statistical fluctuation.

References

- [1] ATLAS Collaboration, Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Phys. Lett. B 716. arXiv:1207.7214, doi:10.1016/j.physletb.2012.08.020.
- [2] CMS Collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, Phys. Lett. B 716. arXiv:1207.7235, doi:10.1016/j.physletb.2012.08.021.

- [3] CMS Collaboration, Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV, JHEP 06 (2013) 081. doi:10.1007/JHEP06(2013)081.
- [4] CMS Collaboration, Study of the Mass and Spin-Parity of the Higgs Boson Candidate via Its Decays to Z Boson Pairs, Phys. Rev. Lett. 110 (2013) 081803. arXiv:1212.6639, doi:10.1103/PhysRevLett.110.081803.
- [5] ATLAS Collaboration, Evidence for the spin-0 nature of the Higgs boson using ATLAS data, Phys. Lett. B 726 (2013) 120. arXiv:1307.1432, doi:10.1016/j.physletb.2013.08.026.
- [6] CMS Collaboration, Evidence for the 125 GeV Higgs boson decaying to a pair of τ leptons, J. High Energy Phys. 05 (2014) 104. doi:10.1007/JHEP05(2014)104.
- [7] A. Dery, A. Efrati, Y. Hochberg, Y. Nir, What if $\text{BR}(H \rightarrow \mu\mu)/\text{BR}(H \rightarrow \tau\tau)$ does not equal m_μ^2/m_τ^2 ?, JHEP 05 (2013) 039. arXiv:1302.3229, doi:10.1007/JHEP05(2013)039.
- [8] CMS Collaboration, Search for Lepton Flavour Violating Decays of the Higgs Boson, CMS Physics Analysis Summary CMS-PAS-HIG-14-005 (2014). URL <http://cds.cern.ch/record/1740976>
- [9] CMS Collaboration, Search for the standard model Higgs boson produced in association with a W or a Z boson and decaying to bottom quarks, Phys.Rev. D89 (1) (2014) 012003. arXiv:1310.3687, doi:10.1103/PhysRevD.89.012003.
- [10] CMS Collaboration, Evidence for the direct decay of the 125 GeV Higgs boson to fermions, Nature Phys. 10. doi:10.1038/nphys3005.
- [11] CMS Collaboration, Search for standard model Higgs boson production in the $\mu^+\mu^-$ final state with the CMS experiment in pp collisions at $\sqrt{s} = 7$ and 8 TeV, CMS Physics Analysis Summary CMS-PAS-HIG-13-007 (2013). URL <http://cdsweb.cern.ch/record/1606831>
- [12] R. Harnik, J. Kopp, J. Zupan, Flavor Violating Higgs Decays, JHEP 03 (2013) 026. arXiv:1209.1397, doi:10.1007/JHEP03(2013)026.