



Probing CP violation in $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ decays

Tim Gershon, Thomas Latham and Rafael Silva Coutinho

Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

Abstract

The three-body charmless hadronic decay $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ provides a number of novel possibilities to search for CP violation effects and test the Standard Model of particle physics. These include fits to the Dalitz-plot distributions of the decay-time-integrated final state, decay-time-dependent (but without initial state flavour tagging) fits to the Dalitz-plot distribution, as well as full decay-time-dependent and flavour tagged fits. The relative sensitivities of these different approaches are investigated.

Keywords: CP violation, b -hadron decays, Dalitz-plot analysis

1. Introduction

The search for a new source of CP violation in addition to that predicted by the CKM matrix [1, 2] is among the main goals of current particle physics research. In the quark sector, a number of important tests have been performed by experiments such as BaBar, Belle and LHCb [3–7]. This line of investigation will be continued by Belle II [8] and the upgraded LHCb experiment [9, 10].

One of the most interesting approaches to search for new sources of CP violation is by studying the decay-time distribution of neutral B meson decays to hadronic final states mediated by the loop (“penguin”) $b \rightarrow s$ amplitude. As-yet undiscovered particles can contribute in the loops and cause the observables to deviate from their expected values in the Standard Model (SM) [11–14]. Studies of B^0 decays to ϕK_s^0 , $\eta' K_s^0$, $K_s^0 K_s^0 K_s^0$ and various other final states have been performed for this reason. The latest results are consistent with the SM predictions, but improved measurements are needed to be sensitive to small deviations.

Experience from previous experiments has shown that full decay-time-dependent Dalitz-plot analysis of a three-body decay (for example $B^0 \rightarrow K_s^0 \pi^+ \pi^-$) is more sensitive than a “quasi-two-body” approach (in this example, considering only the $K_s^0 \rho^0$ contribution).

This is particularly notable in the case that broad resonances contribute, since interference causes effects to which quasi-two-body approaches have no sensitivity [15–17]. Several methods have been proposed to exploit such interferences in $b \rightarrow s$ transitions to allow determination of underlying parameters such as the CKM phase γ with reduced theoretical uncertainty [18–22]. Full decay-time-dependent Dalitz-plot analyses of $B^0 \rightarrow K_s^0 \pi^+ \pi^-$ [23, 24] and $B^0 \rightarrow K_s^0 K^+ K^-$ [25, 26] have been performed by BaBar and Belle, but similar studies of B_s^0 meson decays have not yet been possible.

First results from LHCb on decays of the B_s^0 meson via hadronic $b \rightarrow s$ amplitudes have, however, recently become available. Decay-time-dependent analyses of $B_s^0 \rightarrow K^+ K^-$ [27] and $B_s^0 \rightarrow \phi \phi$ [28] have already been performed. The first observations of $B_s^0 \rightarrow K_s^0 K^\pm \pi^\mp$ and $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ have also been reported [29], including information on contributing K^* resonances [30], suggesting that it will be possible to study CP violation in these modes in the future.

One interesting feature of the $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ decays is that an asymmetry in the time-integrated yields across the mirror line of the Dalitz plot is a signature of CP violation [31–33]. This can be exploited to search for CP asymmetry with either model-independent or model-dependent approaches. Another important aspect of the

B_s^0 system, with regard to CP violation searches, is the non-zero width difference $\Delta\Gamma_s$ between the mass eigenstates. Compared to the situation for B^0 decays, the decay-time distribution receives additional terms that do not vanish when integrated over the initial flavour of the B meson. This implies that information about CP violation parameters can be obtained from analyses that do not tag the initial flavour, through so-called effective lifetime measurements [34, 35]. Although analyses that include flavour tagging information will always be more sensitive, this method may still be of interest for analyses based on small event samples, since it is difficult to achieve high effective tagging efficiency at hadron collider experiments such as LHCb.

The purpose of this paper is to investigate the comparative sensitivity of different methods to search for CP violation in $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ decays. The methods that are considered are (i) untagged, decay-time-integrated; (ii) untagged, decay-time-dependent; (iii) tagged, decay-time-dependent. Only model-dependent methods are included. The study is based on a simple toy model for the decays, including contributions only from $K^*(892)$, $K_0^*(1430)$, $\rho(770)$, and $f_0(980)$ resonances, implemented with the Laura++ Dalitz-plot fitting package [36].

2. Formalism

The decay-time distribution for the decays of mesons, initially produced as \bar{B}_s^0 and B_s^0 flavour eigenstates, to a final state f can be written [37]

$$\frac{d}{dt} \Gamma_{\bar{B}_s^0 \rightarrow f}(t) = \frac{N_f e^{-t/\tau(B_s^0)}}{2\tau(B_s^0)} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + S_f \sin(\Delta m_s t) - C_f \cos(\Delta m_s t) + A_f^{\Delta\Gamma_s} \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right], \quad (1)$$

and

$$\frac{d}{dt} \Gamma_{B_s^0 \rightarrow f}(t) = \frac{N_f e^{-t/\tau(B_s^0)}}{2\tau(B_s^0)} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - S_f \sin(\Delta m_s t) + C_f \cos(\Delta m_s t) + A_f^{\Delta\Gamma_s} \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right], \quad (2)$$

where the mass and width differences between the light (L) and heavy (H) B_s^0 physical eigenstates are defined as $\Delta m_s = m_H - m_L$ and $\Delta\Gamma_s = \Gamma_L - \Gamma_H$, and the B_s^0 lifetime is $\tau(B_s^0) = \left(\frac{\Gamma_L + \Gamma_H}{2}\right)^{-1}$ (units with $\hbar = c = 1$ are used). The coefficients of the $\sin(\Delta m_s t)$, $\cos(\Delta m_s t)$ and $\sinh\left(\frac{\Delta\Gamma_s t}{2}\right)$ terms are often expressed as

$$S_f \equiv \frac{2 \Im(\lambda_f)}{1 + |\lambda_f|^2}, C_f \equiv \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, A_f^{\Delta\Gamma_s} \equiv -\frac{2 \Re(\lambda_f)}{1 + |\lambda_f|^2}, \quad (3)$$

where the parameter λ_f encodes information about CP violation and is given by $\lambda_f = \frac{q}{p} \frac{\bar{\mathcal{A}}_f}{\mathcal{A}_f}$ where $\bar{\mathcal{A}}_f$ and \mathcal{A}_f are the amplitudes for \bar{B}_s^0 and B_s^0 decay to the final state f and q and p define the physical eigenstates in terms of their flavour components

$$|B_{sL}^0\rangle = p|B_s^0\rangle + q|\bar{B}_s^0\rangle, \quad |B_{sH}^0\rangle = p|B_s^0\rangle - q|\bar{B}_s^0\rangle, \quad (4)$$

with $|p|^2 + |q|^2 = 1$. Note that, by definition,

$$(S_f)^2 + (C_f)^2 + (A_f^{\Delta\Gamma_s})^2 = 1. \quad (5)$$

In the remainder of this work, it will be assumed that $|q/p| = 1$ (*i.e.* absence of CP violation in mixing).

By requiring that the integral over t from zero to infinity of the sum of Eq. (1) and Eq. (2) is equal to $|\mathcal{A}_f|^2 + |\bar{\mathcal{A}}_f|^2$, the normalisation factor is found to be

$$\mathcal{N}_f = \left(|\mathcal{A}_f|^2 + |\bar{\mathcal{A}}_f|^2 \right) \frac{1 - y^2}{1 + y A_f^{\Delta\Gamma_s}}, \quad (6)$$

where $y = \tau(B_s^0) \Delta\Gamma_s / 2$. The correction involving y is the origin of the difference between branching fractions calculated at $t = 0$ or after integration over decay time [38].

The discussion above is appropriate for any final state f , including two-body decays. For multibody decays described by the isobar model [39–41], the total amplitude is obtained from a sum of amplitudes from resonant or nonresonant decay channels,

$$\mathcal{A}_f = \sum_{j=1}^N c_j F_j(f), \quad \bar{\mathcal{A}}_f = \sum_{j=1}^N \bar{c}_j F_j(f), \quad (7)$$

where $F_j(f)$ are dynamical amplitudes that contain the lineshape and spin-dependence of the hadronic part of the amplitude labelled by j evaluated at the point in phase space given by f , and c_j are complex coefficients describing the relative magnitude and phase of the different decay channels. Since the $F_j(f)$ terms describe strong dynamics only, they are CP conserving. By contrast, the c_j terms can be CP violating, which is manifested when \bar{c}_j differs from c_j in either magnitude or phase – typically this can occur when the amplitude j has contributions from both “tree” and “loop” (or “penguin”) amplitudes.

The above discussion makes clear how different forms of CP violation may be manifest in different types of analysis:

- i. Untagged, decay-time-integrated Dalitz plot:
In the absence of all forms of CP violation, there is a symmetry between the mirror line in the $K_s^0 \pi^+ \pi^-$

phase-space. This can be broken, for example, by CP violation in decay to flavour-specific final states, such as $K^{\pm}\pi^{\mp}$, since the B_s^0 and \bar{B}_s^0 decays populate different regions of the Dalitz plot. In general one would expect to find larger asymmetries in some local regions of the phase space, and either model-dependent or model-independent methods could be used to search for such effects. A model-dependent fit can determine the C_f parameters of Eq. 1 and 2.

- ii. Untagged, decay-time-dependent Dalitz plot: The $A_f^{\Delta\Gamma_s}$ terms of Eq. 1 and 2 can be determined, and therefore more information is obtained compared to the decay-time-integrated case.
- iii. Tagged, decay-time-dependent Dalitz plot: All terms, including the S_f parameters, can be determined. This method therefore provides additional sensitivity to the model parameters, in particular to the relative phase between B_s^0 and \bar{B}_s^0 decay amplitudes.

This general discussion does not answer the question of how much additional sensitivity is obtained as the analysis is made increasingly more complex. That will be addressed in the next sections.

3. Method to generate toy samples

Several ensembles of Monte Carlo pseudoexperiments are generated to investigate CP violation effects in $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ decays. The simulation is performed without any experimental effects, such as background, acceptance, resolution or imperfect flavour tagging. The toy model contains the $\rho^0(770)$, $f_0(980)$, $K^{*\pm}(892)$ and $K_0^{*\pm}(1430)$ resonances. All mass terms are described by the relativistic Breit-Wigner (RBW) function, apart from the $K_0^{*\pm}(1430)$ lineshape which is modelled by the LASS shape [42]. The parametrisation of complex coefficients is given by

$$\bar{c}_j^{(-)} = (x_j \pm \Delta x_j) + i(y_j \pm \Delta y_j), \quad (8)$$

where Δx_j and Δy_j are CP -violating parameters. Table 1 summarises the baseline model used to generate events, with decay-time distribution given in Eq. 1 and 2.

In the fit, the $\bar{c}_j^{(-)}$ coefficients are measured relative to the $\rho^0(770)$ resonance contribution. Each pseudoexperiment is fitted many times with randomised initial values of the parameters in order to find the global minimum of the negative log likelihood function. Asymmetries are calculated as

$$\mathcal{A}_{CPj} = \frac{|\bar{c}_j^{(-)}|^2 - |c_j|^2}{|\bar{c}_j^{(-)}|^2 + |c_j|^2} = \frac{-2(x_j \Delta x_j + y_j \Delta y_j)}{x_j^2 + \Delta x_j^2 + y_j^2 + \Delta y_j^2}. \quad (9)$$

Table 1: Benchmark parameters for the baseline Dalitz plot model used as input in the generation.

Resonance	x_j	Δx_j	y_j	Δy_j
$\rho^0(770)$	1.0	0.0	0.0	0.0
$f_0(980)$	$0.4 \cos(5\pi/4)$	0.0	$0.4 \sin(5\pi/3)$	0.0
$K^{*\pm}(892)$	$1.2 \cos(\pi/3)$	0.0	$1.2 \sin(\pi/3)$	0.0
$K_0^{*\pm}(1430)$	$1.7 \cos(\pi/3)$	0.0	$1.7 \sin(\pi/3)$	0.0

CP violation can also be manifest in a difference between the phase of the B_s^0 and \bar{B}_s^0 decay amplitudes,

$$\Delta\delta_j = \arg\left(\frac{\bar{c}_j}{c_j}\right) = \tan^{-1}\left(\frac{y_j + \Delta y_j}{x_j + \Delta x_j}\right) - \tan^{-1}\left(\frac{y_j - \Delta y_j}{x_j - \Delta x_j}\right). \quad (10)$$

The baseline model is modified in various ways to introduce CP violation. Interference between the B_s^0 - \bar{B}_s^0 oscillation and decay amplitudes is incorporated through the CP violation weak phase ϕ_s . While the SM predicts $\phi_s^{\text{SM}} = -2\beta_s \equiv -2 \arg(-V_{ts} V_{tb}^* / V_{cs} V_{cb}^*) = -0.036 \pm 0.002$ rad, contributions from physics beyond the SM could lead to much larger values. Three different scenarios are generated: $\phi_s = 0$, $\phi_s = -2\beta_s$ and $\phi_s = -20\beta_s$. In addition, CP violation in the decay of each resonance is examined: CP violation in the magnitude, with $\mathcal{A}_{CP} = 5\%$, 10% , 20% and 50% ; CP violation due to the difference in the relative phase in steps of $\pi/4$ from 0 to 2π ; and CP violation in both magnitude and phase difference. Pseudoexperiments are generated with sample size corresponding roughly to the anticipated yields available at LHCb by the end of the LHC Run II (2000 events). Ensembles with other sample sizes are also generated to test the scaling of the uncertainties. Only a representative subset of the results obtained are presented here due to space constraints.

4. Results

Figure 1 shows the results for various scenarios of CP violation in the $K^{*\pm}(892)$ amplitude, with yields corresponding to LHC Run I+II. The fitted values of the isobar coefficients in each pseudoexperiment are represented by the points in the Argand plane, with the ellipses illustrating the central values and 1σ contour boundaries from the ensemble. The colour schemes for B_s^0 and \bar{B}_s^0 coefficients are represented respectively by: blue and cyan for method i; light and dark green for method ii; and red and magenta for method iii.

It is immediately clear that the magnitude of the amplitude is determined much more precisely than the




Figure 1: Fitted values of the $K^{*\pm}(892)$ isobar coefficients plotted in the Argand plane. The points are the values determined from individual pseudoexperiments while the ellipses illustrate the mean value and 68% confidence level contour from the corresponding ensemble. The results for $c_j(\bar{c}_j)$ are shown for method i in blue (cyan), for method ii in green (dark green) and for method iii in red (magenta). All experiments are generated with 2000 signal events and $\phi_s = -2\beta_s$ and the following scenarios: (top left) no CP violation, (top right) $\mathcal{A}_{CP} = 50\%$, (bottom left) $\Delta\delta = 3\pi/4$ and (bottom right) $\mathcal{A}_{CP} = 50\%$ and $\Delta\delta = 3\pi/4$.

phase, leading to the arc-like distribution of points. Table 2 compares the precision of the different fitting methods for each of the CP violation scenarios. The results indicate that the generated asymmetries are retrieved in all scenarios with good precision and without significant bias. The untagged methods give statistical uncertainties that are only slightly larger, due to the fact that the K^* resonances from the decay of B_s^0 and \bar{B}_s^0 populate different regions of the Dalitz plot. In addition, the very similar uncertainties given by the two untagged approaches suggests that the $A_f^{\Delta\Gamma_s}$ term does not provide a significant amount of extra sensitivity. Further studies with realistic experimental effects are necessary to determine the exact sensitivities achievable. An extrapolation of the precision estimated here suggests that such measurements appear to be feasible, albeit with large uncertainty, with the LHCb Run I dataset that is already in hand.

A further study is performed to investigate the sensitivity to the ϕ_s observable. Figure 2 compares the results from methods ii and iii (such a determination is not possible with method i). There is a clear improvement in the determination of the weak phase when tagging is applied. With perfect tagging, the precision on ϕ_s shows

an order of magnitude improvement. Using a more realistic tagging power of $\sim 5\%$, as achieved recently by LHCb [43, 44], still provides a factor ~ 2.5 better sensitivity to ϕ_s than the untagged case. Alternatively one can fix the value of $\phi_s = -2\beta_s$ in the fit and float the Δy_j parameter of the $\rho^0(770)$ resonance in order to measure the relative phase between the B_s^0 and \bar{B}_s^0 decay to this state. This approach is also illustrated in Fig. 2 and shows the same behaviour comparing methods ii and iii.

5. Summary

The recent observation of B_s^0 decays to charmless three-body final states marks the start of a new and interesting field of CP violation investigation. In this note, a comparative sensitivity study for different approaches to Dalitz plot analysis has been performed for $B_s^0 \rightarrow K_s^0 \pi^+ \pi^-$ decays. It has been demonstrated that good precision for the phase difference between B_s^0 and \bar{B}_s^0 decays to $K^{*\pm}(892)\pi^\mp$ can be achieved with untagged analysis approaches (e.g. for the LHC Run I and II). Flavour tagging is, however, needed to determine ϕ_s (i.e. the relative phase in $B_s^0(\bar{B}_s^0) \rightarrow K_s^0 \rho^0(770)$ decays).

Table 2: Comparison of the uncertainties on the $K^{*\pm}(892)$ CP -violating parameters determined using the different fitting methods. The results are quoted in terms of the polar co-ordinates $c_j = a_j e^{i\delta_j}$, $\bar{c}_j = \bar{a}_j e^{i\bar{\delta}_j}$. The relative uncertainties for method iii are quoted (the central values of the parameters correspond to the values given in Table 1, modified according to the CP violation parameters), together with comparisons of the uncertainties with the different Dalitz plot fit methods. The typical uncertainty on the relative precision is ± 0.1 .

CP -violation parameters			$\sigma(\text{method iii})$ (%)				$\frac{\sigma(\text{method ii})}{\sigma(\text{method iii})}$				$\frac{\sigma(\text{method i})}{\sigma(\text{method iii})}$			
\mathcal{A}_{CP}	$\Delta\delta$	ϕ_s	a_j	\bar{a}_j	δ_j	$\bar{\delta}_j$	a_j	\bar{a}_j	δ_j	$\bar{\delta}_j$	a_j	\bar{a}_j	δ_j	$\bar{\delta}_j$
20%	0	0	4.6	3.7	12.3	11.6	1.1	1.2	1.8	1.8	1.0	1.0	1.7	1.8
50%	0	$2\beta_s$	5.1	3.3	15.0	12.2	1.0	1.1	1.6	1.8	0.9	1.1	1.5	1.8
0%	$\pi/4$	0	4.2	4.3	12.2	7.7	1.0	1.1	1.8	1.6	1.0	0.9	1.6	1.5
0%	$3\pi/4$	$2\beta_s$	4.2	4.0	12.4	4.7	1.0	1.2	1.6	1.8	1.0	1.1	1.5	1.8
5%	$\pi/4$	0	4.5	3.9	11.4	8.3	1.0	1.1	1.8	1.4	0.9	1.0	1.8	1.5
50%	π	$2\beta_s$	5.2	3.6	14.5	7.1	1.0	1.1	1.6	1.7	1.1	0.9	1.6	1.7

These results indicate directions for possible amplitude analyses that can be pursued in future by Belle II and LHCb.

Acknowledgements

This work is funded by the European Research Council under FP7 and by the United Kingdom's Science and Technology Facilities Council. The authors thank their colleagues in the LHCb collaboration for the experimental progress that provides the context for this work.

References

[1] N. Cabibbo, Unitary symmetry and leptonic decays, Phys.Rev.Lett. 10 (1963) 531. doi:10.1103/PhysRevLett.10.531.

[2] M. Kobayashi, T. Maskawa, CP violation in the renormalizable theory of weak interaction, Prog.Theor.Phys. 49 (1973) 652. doi:10.1143/PTP.49.652.

[3] M. Antonelli, et al., Flavor physics in the quark sector, Phys.Rept. 494 (2010) 197. arXiv:0907.5386, doi:10.1016/j.physrep.2010.05.003.

[4] J. Brodzicka, et al., Physics achievements from the Belle experiment, PTEP 2012 (2012) 04D001. arXiv:1212.5342, doi:10.1093/ptep/pts072.

[5] A. Bevan, et al., The physics of the B factories arXiv:1406.6311.

[6] R. Aaij et al., and A. Bharucha, et al., Implications of LHCb measurements and future prospects, Eur. Phys. J. C73 (2013) 2373. arXiv:1208.3355, doi:10.1140/epjc/s10052-013-2373-2.

[7] Y. Amhis, et al., Averages of b -hadron, c -hadron, and τ -lepton properties as of early 2012, updated results and plots available at: <http://www.slac.stanford.edu/xorg/hfag/>. arXiv:1207.1158.

[8] T. Aushev, et al., Physics at Super B Factory arXiv:1002.5012.

[9] LHCb collaboration, Letter of Intent for the LHCb Upgrade (CERN-LHCC-2011-001. LHCC-I-018).

[10] LHCb collaboration, Framework TDR for the LHCb Upgrade (CERN-LHCC-2012-007. LHCB-TDR-012).

[11] Y. Grossman, M. P. Worah, CP asymmetries in B decays with new physics in decay amplitudes, Phys. Lett. B395 (1997) 241. arXiv:hep-ph/9612269, doi:10.1016/S0370-2693(97)00068-3.

[12] R. Fleischer, CP violation and the role of electroweak penguins in nonleptonic B decays, Int. J. Mod. Phys. A12 (1997) 2459. arXiv:hep-ph/9612446, doi:10.1142/S0217751X97001432.

[13] D. London, A. Soni, Measuring the CP angle β in hadronic $b \rightarrow s$ penguin decays, Phys. Lett. B407 (1997) 61. arXiv:hep-ph/9704277, doi:10.1016/S0370-2693(97)00695-3.

[14] M. Ciuchini, E. Franco, G. Martinelli, A. Masiero, L. Silvestrini, CP violating B decays in the Standard Model and supersymmetry, Phys. Rev. Lett. 79 (1997) 978. arXiv:hep-ph/9704274, doi:10.1103/PhysRevLett.79.978.

[15] A. E. Snyder, H. R. Quinn, Measuring CP asymmetry in $B \rightarrow \rho\pi$ decays without ambiguities, Phys.Rev. D48 (1993) 2139. doi:10.1103/PhysRevD.48.2139.

[16] J. Charles, A. Le Yaouanc, L. Oliver, O. Pene, J. C. Raynal, $B^0(t) \rightarrow DPP$ time-dependent Dalitz plots, CP-violating angles 2β , $2\beta + \gamma$, and discrete ambiguities, Phys. Lett. B425 (1998) 375. arXiv:hep-ph/9801363, doi:10.1016/S0370-2693(98)00250-0.

[17] T. Latham, T. Gershon, A method to measure $\cos(2\beta)$ using time-dependent Dalitz plot analysis of $B^0 \rightarrow D_{CP}\pi^+\pi^-$, J.Phys. G36 (2009) 025006. arXiv:0809.0872, doi:10.1088/0954-3899/36/2/025006.

[18] M. Ciuchini, M. Pierini, L. Silvestrini, New bounds on the CKM matrix from $B \rightarrow K\pi\pi$ Dalitz plot analyses, Phys.Rev. D74 (2006) 051301. arXiv:hep-ph/0601233, doi:10.1103/PhysRevD.74.051301.

[19] M. Ciuchini, M. Pierini, L. Silvestrini, Hunting the CKM weak phase with time-integrated Dalitz analyses of $B_s^0 \rightarrow KK\pi$ and $B_s^0 \rightarrow K\pi\pi$ decays, Phys.Lett. B645 (2007) 201. arXiv:hep-ph/0602207, doi:10.1016/j.physletb.2006.12.043.

[20] M. Gronau, D. Pirjol, A. Soni, J. Zupan, Improved method for CKM constraints in charmless three-body B and B_s^0 decays, Phys.Rev. D75 (2007) 014002. arXiv:hep-ph/0608243, doi:10.1103/PhysRevD.75.014002.

[21] M. Gronau, D. Pirjol, A. Soni, J. Zupan, Constraint on $\bar{\rho}, \bar{\eta}$ from $B \rightarrow K^*\pi$, Phys.Rev. D77 (2008) 057504. arXiv:0712.3751, doi:10.1103/PhysRevD.77.057504.

[22] I. Bediaga, G. Guerrer, J. M. de Miranda, Extracting the quark mixing phase γ from $B^\pm \rightarrow K^\pm\pi^+\pi^-$, $B^0 \rightarrow K_S\pi^+\pi^-$, and $\bar{B}^0 \rightarrow K_S\pi^+\pi^-$, Phys.Rev. D76 (2007) 073011. arXiv:hep-ph/0608268, doi:10.1103/PhysRevD.76.073011.

[23] B. Aubert, et al., Time-dependent amplitude analysis of $B^0 \rightarrow K_S^0\pi^+\pi^-$, Phys. Rev. D80 (2009) 112001. arXiv:0905.3615, doi:10.1103/PhysRevD.80.112001.

[24] J. Dalseno, et al., Time-dependent Dalitz Plot measurement of CP parameters in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays, Phys. Rev. D79 (2009) 072004. arXiv:0811.3665, doi:10.1103/PhysRevD.79.072004.




Figure 2: Fitted values of ϕ_s for (top left) method ii with $\phi_s = -2\beta_s$, (top right) method ii with $\phi_s = -20\beta_s$ and (bottom left) method iii with $\phi_s = -20\beta_s$. The (bottom right) Argand plot displays the fitted coefficient values for the $\rho^0(770)$ resonance with fixed $\phi_s = -2\beta_s$ and $\mathcal{A}_{CP} = 50\%$.

- [25] Y. Nakahama, et al., Measurement of CP violating asymmetries in $B^0 \rightarrow K^+K^-K_S^0$ decays with a time-dependent Dalitz approach, Phys. Rev. D82 (2010) 073011. arXiv:1007.3848, doi:10.1103/PhysRevD.82.073011.
- [26] J. P. Lees, et al., Study of CP violation in Dalitz-plot analyses of $B^0 \rightarrow K^+K^-K_S^0$, $B^+ \rightarrow K^+K^-K^+$, and $B^+ \rightarrow K_S^0K_S^0K^+$, Phys.Rev. D85 (2012) 112010. arXiv:1201.5897, doi:10.1103/PhysRevD.85.112010.
- [27] R. Aaij, et al., First measurement of time-dependent CP violation in $B_s^0 \rightarrow K^+K^-$ decays, JHEP 10 (2013) 183. arXiv:1308.1428, doi:10.1007/JHEP10(2013)183.
- [28] R. Aaij, et al., First measurement of the CP -violating phase in $B_s^0 \rightarrow \phi\phi$ decays, Phys. Rev. Lett. 110 (2013) 241802. arXiv:1303.7125, doi:10.1103/PhysRevLett.110.241802.
- [29] R. Aaij, et al., Study of $B_{(s)}^0 \rightarrow K_S^0 h^+ h^-$ decays with first observation of $B_s^0 \rightarrow K_S^0 K^\pm \pi^\mp$ and $B_s^0 \rightarrow K_S^0 \pi^+ \pi^-$, JHEP 10 (2013) 143. arXiv:1307.7648, doi:10.1007/JHEP10(2013)143.
- [30] R. Aaij, et al., Observation of $B_s^0 \rightarrow K^{*\pm} K^\mp$ and evidence of $B_s^0 \rightarrow K^{*-} \pi^+$ decays, to appear in New J. Phys. arXiv:1407.7704.
- [31] G. Burdman, J. F. Donoghue, B meson CP violation without flavor identification, Phys.Rev. D45 (1992) 187. doi:10.1103/PhysRevD.45.187.
- [32] S. Gardner, Direct CP violation in untagged B meson decays, Phys.Lett. B553 (2003) 261. arXiv:hep-ph/0203152, doi:10.1016/S0370-2693(02)03263-X.
- [33] S. Gardner, J. Tandan, Observing direct CP violation in untagged B meson decays, Phys.Rev. D69 (2004) 034011. arXiv:hep-ph/0308228, doi:10.1103/PhysRevD.69.034011.
- [34] I. Dunietz, B_s^0 - \bar{B}_s^0 mixing, CP violation and extraction of CKM phases from untagged B_s^0 data samples, Phys.Rev. D52 (1995) 3048. arXiv:hep-ph/9501287, doi:10.1103/PhysRevD.52.3048.
- [35] R. Fleischer, R. Knegjens, Effective lifetimes of B_s^0 decays and their constraints on the B_s^0 - \bar{B}_s^0 mixing parameters, Eur.Phys.J. C71 (2011) 1789. arXiv:1109.5115, doi:10.1140/epjc/s10052-011-1789-9.
- [36] Laura++ Dalitz plot fitting package, University of Warwick, <http://laura.hepforge.org/>.
- [37] I. Dunietz, R. Fleischer, U. Nierste, In pursuit of new physics with B_s^0 decays, Phys.Rev. D63 (2001) 114015. arXiv:hep-ph/0012219, doi:10.1103/PhysRevD.63.114015.
- [38] K. De Bruyn, et al., Branching ratio measurements of B_s^0 decays, Phys.Rev. D86 (2012) 014027. arXiv:1204.1735, doi:10.1103/PhysRevD.86.014027.
- [39] G. N. Fleming, Recoupling effects in the isobar model. I. General formalism for three-pion scattering, Phys.Rev. 135 (1964) B551. doi:10.1103/PhysRev.135.B551.
- [40] D. Morgan, Phenomenological analysis of $I = 1/2$ single-pion production processes in the energy range 500 to 700 MeV, Phys.Rev. 166 (1968) 1731. doi:10.1103/PhysRev.166.1731.
- [41] D. Herndon, P. Soding, R. Cashmore, A generalised isobar model formalism, Phys.Rev. D11 (1975) 3165. doi:10.1103/PhysRevD.11.3165.
- [42] D. Aston, et al., A study of $K^- \pi^+$ scattering in the reaction $K^- p \rightarrow K^- \pi^+ n$ at 11-GeV/c, Nucl.Phys. B296 (1988) 493. doi:10.1016/0550-3213(88)90028-4.
- [43] R. Aaij, et al., Measurement of CP asymmetry in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, to appear in JHEP. arXiv:1407.6127.
- [44] R. Aaij, et al., Measurement of the CP -violating phase ϕ_s in $\bar{B}_s^0 \rightarrow D_s^+ D_s^-$ decays, to appear in Phys. Rev. Lett. arXiv:1409.4619.