

Neutron deficient exotic nuclei and the Physics of the "proton rich side" of the nuclear chart



Contribution ID : 4

Type : **not specified**

Studies of Two Proton radioactivity

Monday, 21 February 2011 11:00 (40)

Extremely proton-rich nuclei with odd or even atomic numbers were predicted by V. Goldansky in 1960 to decay through one- or two-proton radioactivity, respectively. Two-proton ($2p$) radioactivity, a spontaneous decay of an atomic nucleus by emission of two protons, is the most recently discovered nuclear disintegration mode. It has first been reported for ^{45}Fe in 2002 with a half-life of about 4 ms, which is 1000 times longer than the quasi-classical estimate of "di-proton" (or 2He) cluster emission. Further observations of $2p$ radioactivity in ^{19}Mg , ^{54}Zn , ^{48}Ni , ^{94}mAg have confirmed unexpectedly large half-lives of $2p$ precursors thus indicating regular existence of long-lived nuclei beyond the proton drip line. The experimental methods used to produce, identify and detect new nuclear species via their $2p$ decay will be reviewed. The up-to-date theoretical understanding of specific observables accessible in the $2p$ decay measurements will be discussed with an emphasis on $2p$ -precursor's nuclear structure.

Summary

Primary author(s) : MUKHA, Ivan (Helholzzentrum Gesellschaft fur SchwerIonenforschung (GSI))

Presenter(s) : MUKHA, Ivan (Helholzzentrum Gesellschaft fur SchwerIonenforschung (GSI))

Session Classification : Monday February 21st, 2011. 11:00 - 13:00