

Shedding light on Dark Matter from accelerator physics

Bryan Zaldívar Montero


CPAN III (2011), Barcelona.

based on Y. Mambrini, B.Z., hep-ph/1106.4819
JCAP 1110 (2011) 023

- Introduction
 - Experimental evidences
 - DM interpretations
 - Nature of couplings
 - Motivation
- Discussion & Results
 - LEP vs Tevatron
 - fermionic DM
 - scalar DM
 - combined analysis (astro+accel)
 - universal lepton coupling
 - pure electron coupling
 - words on LHC
- Conclusions

DARK EXCESSES? (ACC. EXP., D.D)


CDF Collaboration hep-ex/1104.06991




DARK EXCESSES? (ACC. EXP., D.D)

Many possibilities:

- Standard Model effect [Sullivan et al, hep-ph/1104.3790](#)
- Stringy origin [Anchordoqui et al, hep-ph/1104.2303](#)
- Technicolor [Eichten et al, hep-ph/1104.0976](#)
- 2-Higgs models [Chen et al, hep-ph/1105.2870](#)
- etc [hep-ph/11xx.xxx](#)
- Dark Matter interpretation [e.g. Buckley et al, hep-ph/1104.3145](#)
 - "dark" Z-boson
 - χ -nucleon scattering $\sigma_{\chi-n} \sim 10^{-40} \text{cm}^2$ (for $m_\chi \sim 10 \text{GeV}$)
 - **Hadrophilic coupling**




DARK EXCESSES? (ASTRO. EXP., D.D.)



DARK EXCESSES? (ASTRO. EXP., D.D.)


Tension in XENON and DAMA results

- **electrophilic DM** Bernabei et al, astro-ph/0712.0562
DAMA sensible also to electron recoil ($\pm 33\%$)
- **leptophilic** Fox et al, hep-ph/0811.0399
- **hadrophilic** Kopp et al, hep-ph/0907.3159




DARK EXCESSES? (ASTRO. EXP. I.D.)

- γ -rays observed by Fermi
- inner parsecs unexplained
- **leptophilic DM**






DARK EXCESSES? (ASTRO. EXP. I.D.)



- e^+ -flux, by PAMELA
- electrophilic, μ -philic, π -philic, ...



Cholis et al, astro-ph/0810.5344





MOTIVATION (LEP vs. TEVATRON)




- $\mathcal{O}_V = \frac{(\bar{\chi}\gamma_\mu\chi)(\bar{l}\gamma^\mu l)}{\Lambda^2}$
- $\mathcal{O}_S = \frac{(\bar{\chi}\chi)(\bar{l}l)}{\Lambda^2}$
- $\mathcal{O}_A = \frac{(\bar{\chi}\gamma_\mu\gamma_5\chi)(\bar{l}\gamma^\mu\gamma_5 l)}{\Lambda^2}$
- $\mathcal{O}_t = \frac{(\bar{\chi}l)(\bar{l}\chi)}{\Lambda^2}$

MOTIVATION (LEP vs. TEVATRON)




- $\mathcal{O}_V = \frac{(\bar{\chi} \gamma_\mu \chi)(\bar{l} \gamma^\mu l)}{\Lambda^2}$
- $\mathcal{O}_S = \frac{(\bar{\chi} \chi)(\bar{l} l)}{\Lambda^2}$
- $\mathcal{O}_A = \frac{(\bar{\chi} \gamma_\mu \gamma_5 \chi)(\bar{l} \gamma^\mu \gamma_5 l)}{\Lambda^2}$
- $\mathcal{O}_t = \frac{(\bar{\chi} l)(\bar{l} \chi)}{\Lambda^2}$




MOTIVATION (LEP vs. TEVATRON)

$$\bullet \sigma_{1\text{-loop}} = \frac{4\alpha^2 \mu_p^2}{18^2 \pi^3 \Lambda^4} \cdot [\sum_l f(q^2, m_l)]^2$$




$$\bullet (\sigma v)_s = \frac{1}{8\pi\Lambda^4} \sqrt{1 - \frac{m_\tau^2}{m_\chi^2} (m_\chi^2 - m_l^2)} v_{\text{rel}}^2$$

MOTIVATION (LEP vs. TEVATRON)





$$\bullet \sigma_{1\text{-loop}} = \frac{4\alpha^2 \mu_p^2}{18^2 \pi^3 \Lambda^4} \cdot [\sum_l f(q^2, m_l)]^2$$

C





$$\bullet (\sigma v)_S = \frac{1}{8\pi \Lambda^4} \sqrt{1 - \frac{m_l^2}{m_\chi^2}} (m_\chi^2 - m_l^2) v_{\text{rel}}^2$$

MOTIVATION (LEP vs. TEVATRON)





- Luminosity of 1 fb^{-1}


MOTIVATION (LEP vs. TEVATRON)




- Luminosity of 1 fb^{-1}




MOTIVATION (LEP vs. TEVATRON)




MOTIVATION (LEP vs. TEVATRON)




MOTIVATION (LEP vs. TEVATRON)



MOTIVATION (LEP vs. TEVATRON)



MOTIVATION (LEP vs. TEVATRON)



THE MODELS

- Effective scales: $\frac{1}{\Lambda_l} \equiv \frac{\sqrt{g_l}}{\Lambda}$; $\frac{1}{\Lambda_h} \equiv \frac{\sqrt{g_h}}{\Lambda}$

- Operators:

$$\mathcal{L}_V = \sum_i \frac{g_l^i}{\Lambda^2} (\bar{l}^i \gamma^\mu l^i) (\bar{\chi} \gamma_\mu \chi) + \sum_i \frac{g_h^i}{\Lambda^2} (\bar{q}^i \gamma^\mu q^i) (\bar{\chi} \gamma_\mu \chi)$$

$$\mathcal{L}_S = \sum_i \frac{g_l^i}{\Lambda^2} (\bar{l}^i l^i) (\bar{\chi} \chi) + \sum_i \frac{g_h^i}{\Lambda^2} (\bar{q}^i q^i) (\bar{\chi} \chi)$$

$$\mathcal{L}_A = \sum_i \frac{g_l^i}{\Lambda^2} (\bar{l}^i \gamma^\mu \gamma^5 l^i) (\bar{\chi} \gamma_\mu \gamma^5 \chi) + \sum_i \frac{g_h^i}{\Lambda^2} (\bar{q}^i \gamma^\mu \gamma^5 q^i) (\bar{\chi} \gamma_\mu \gamma^5 \chi)$$

$$\mathcal{L}_t = \sum_i \frac{g_l^i}{\Lambda^2} (\bar{l}^i \chi) (\bar{\chi} l^i) + \sum_i \frac{g_h^i}{\Lambda^2} (\bar{q}^i \chi) (\bar{\chi} q^i)$$

- Models in lepton sector:

A) Electrophilic couplings: $g_l^e = g_e$, $g_l^{i=\mu,\tau,\nu_i} = 0$

B) Charged lepton couplings: $g_l^{i=e,\mu,\tau} = g_l$, $g_l^{i=\nu_i} = 0$

C) Universal lepton couplings: $g_l^{i=e,\mu,\tau,\nu_i} = g_l$

- Universality in hadronic couplings: $g_h^{i=u,d,c,s,b,t} = g_h$.

ANNIHILATION CROSS-SECTIONS

$$\frac{d\sigma_I}{d\Omega} = \frac{|\mathcal{M}_I|^2}{64\pi^2 s} \frac{\sqrt{s - 2m_3^2 - 2m_4^2 + \frac{(m_3^2 - m_4^2)^2}{s}}}{\sqrt{s - 4m_\chi^2}}; \quad s \simeq 4m_\chi^4 + m_\chi^2 v^2$$

$$\sigma_I^J v = g_l^2 \sum_{l=e,\mu,\tau,\nu} \sigma_{I,l}^J v + c g_h^2 \sum_{h=u,d,c,s,t,b} \sigma_{I,h}^J v$$

J : operator type
 I : coupling type

$$\sigma_{V,k} v = 4g_\Lambda \left(24(2m_\chi^2 + m_k^2) + \frac{8m_\chi^4 - 4m_\chi^2 m_k^2 + 5m_k^4}{m_\chi^2 - m_l^2} v^2 \right)$$

$$\sigma_{S,k} v = 24g_\Lambda (m_\chi^2 - m_k^2) v^2$$

$$g_\Lambda = \frac{\sqrt{1 - m_k^2/m_\chi^2}}{192\pi\Lambda^4}$$

$$\sigma_{A,k} v = 4g_\Lambda \left(24m_k^2 + \frac{8m_\chi^4 - 22m_\chi^2 m_k^2 + 17m_k^4}{m_\chi^2 - m_k^2} v^2 \right)$$

Requiring




$$\sigma_l^{max} v + \sigma_h^{max} v \gtrsim 3 \times 10^{-26} \text{cm}^3 \text{s}^{-1} \simeq 2.5 \times 10^{-9} \text{GeV}^{-2}$$

assume e.g. $m_\chi \sim 5 \text{ GeV}$, in a model with only **electronic** coupling and **vector-like** interaction:

- $(\Lambda_e)_{\min} \simeq 480 \text{ GeV}$ from LEP
- $m_\chi \gg m_h, m_l$
- $\sigma_V v \simeq \frac{m_\chi^2}{\pi \Lambda_e^4} + 3 \frac{m_\chi^2}{\pi \Lambda_h^4} = \frac{m_\chi^2}{\pi \Lambda_e^4} (1 + 3 \frac{g_h^2}{g_l^2}) \gtrsim 2.5 \times 10^{-9}$
 $\Rightarrow \frac{g_h}{g_l} \simeq 16 \quad 94\% \text{ ann. rate to hadrons}$


NUMERICAL RESULTS

- from LEP bounds:




- Most-conservative: Universal-leptonic, vector
- Less-conservative: Electronic, scalar





COMBINED ANALYSIS





COMBINED ANALYSIS



NUMERICAL RESULTS: UNIVERSAL-LEPTONIC, VECTOR



NUMERICAL RESULTS: ELECTRONIC, SCALAR



SCALAR DARK MATTER

- $\mathcal{L}_e = \frac{g_e}{\Lambda_S} \chi \chi \bar{e} e$
- Madgraph analysis:
 - same bckgr: $e^+ e^- \rightarrow \gamma \nu \bar{\nu}$
 - χ^2 -analysis
 - signal+bckgr sim.
- Infer bounds for $\langle \sigma_{S,e}^s v \rangle$:



$$\sigma_{S,e}^s v \simeq$$

$$\frac{g_e^2}{4\pi\Lambda_S^2} \left(1 - \frac{m_e^2}{m_\chi^2}\right)^{3/2} + \frac{g_e^2}{32\pi\Lambda_S^2} v^2.$$



Result:

- $\sigma_{S,e}^s \simeq 10^{-24} \text{cm}^3/\text{s}$
- **No constraints from LEP, nor Tevatron**
- $\frac{\Lambda_S}{g_e} \Big|_{\langle \sigma v \rangle \sim 10^{-26}} \gtrsim 5 \text{TeV}$



LHC?




LHC (FOX ET AL. PH/1109.4398)



ATLAS 7TeV, 1fb⁻¹ VeryHighPt



Annihilation into $\bar{q}q$





CONCLUSIONS

- Nature of DM coupling is crucial to fit all present data
- We computed the rate of hadronic/leptonic coupling to respect:
 - ① LEP + Tevatron (mono-photon, mono-jet events)
 - ② WMAP + XENON100
- A very light **fermionic DM** ($\lesssim 10$ GeV) **mainly excluded** whatever the type of interaction
- Heavier candidates ($\gtrsim 10$ GeV) should be **largely hadrophobic** (vector int.) or even **excluded** (scalar int.)
- Models with electrophilic couplings (motivated by INTEGRAL, or Synchrotron radiation data) are excluded by LEP/Tevatron analysis

Escaping conclusions: DM candidate not coupled to electrons (LEP bound not applicable) or hadronic coupling only to bottom or charm (Tevatron bound not applicable)

- Nothing to say for the moment for scalar DM.

BACK-UP. LIGHT-MEDIATORS



BACK-UP. HIGH-ENERGY COMPLETIONS

