


AI meeting




M. González-Alonso

- Physics problem: Model-independent searches of New Physics effects (EFT approach) in flavour / LHC / neutrino / nuclear data.
- Minimisation of a likelihood that depends (nonlinearly) on a large number of parameters (Wilson coefficients, hadronic parameters, nuisance parameters, ...).
- SMEFT at the LHC (Brehmer et al'18, Freitas et al'19, ...)

Example: Electroweak precision measurements in the SMEFT [Falkowski, MGA & Mimouni, 2017]



W,Z

$\begin{pmatrix} \delta g_L^{We} \\ \delta g_L^{W\mu} \\ \delta g_L^{W\tau} \\ \delta g_L^{Ze} \\ \delta g_L^{Z\mu} \\ \delta g_L^{Z\tau} \\ \delta g_R^{Ze} \\ \delta g_R^{Z\mu} \\ \delta g_R^{Z\tau} \\ \delta g_L^{Zu} \\ \delta g_L^{Zc} \\ \delta g_L^{Zt} \\ \delta g_R^{Zu} \\ \delta g_R^{Zc} \\ \delta g_L^{Zd} \\ \delta g_L^{Zs} \\ \delta g_R^{Zd} \\ \delta g_R^{Zs} \\ \delta g_R^{Zb} \\ \delta g_L^{Wq_1} \\ \delta g_R^{Wq_1} \end{pmatrix} = \begin{pmatrix} -1.00 \pm 0.64 \\ -1.36 \pm 0.59 \\ 1.95 \pm 0.79 \\ -0.023 \pm 0.028 \\ 0.01 \pm 0.12 \\ 0.018 \pm 0.059 \\ -0.033 \pm 0.027 \\ 0.00 \pm 0.14 \\ 0.042 \pm 0.062 \\ -0.8 \pm 3.1 \\ -0.15 \pm 0.36 \\ -0.3 \pm 3.8 \\ 1.4 \pm 5.1 \\ -0.35 \pm 0.53 \\ -0.9 \pm 4.4 \\ 0.9 \pm 2.8 \\ 0.33 \pm 0.17 \\ 3 \pm 16 \\ 3.4 \pm 4.9 \\ 2.30 \pm 0.88 \\ -1.3 \pm 1.7 \end{pmatrix} \times 10^{-2}$	 $\begin{pmatrix} [c_{\ell\ell}]_{1111} \\ [c_{\ell\ell}]_{1111} \\ [c_{ee}]_{1111} \\ [c_{\ell\ell}]_{1221} \\ [c_{\ell\ell}]_{1122} \\ [c_{\ell\ell}]_{1122} \\ [c_{\ell\ell}]_{2211} \\ [c_{ee}]_{1122} \\ [c_{\ell\ell}]_{1331} \\ [c_{\ell\ell}]_{1133} \\ [c_{\ell\ell}]_{1133} \\ [c_{\ell\ell}]_{3311} \\ [c_{ee}]_{1133} \\ [\hat{c}_{\ell\ell}]_{2222} \\ [c_{\ell\ell}]_{2332} \end{pmatrix} = \begin{pmatrix} 1.01 \pm 0.38 \\ -0.22 \pm 0.22 \\ 0.20 \pm 0.38 \\ -4.8 \pm 1.6 \\ 1.5 \pm 2.1 \\ 1.5 \pm 2.2 \\ -1.4 \pm 2.2 \\ 3.4 \pm 2.6 \\ 1.5 \pm 1.3 \\ 0 \pm 11 \\ -2.3 \pm 7.2 \\ 1.7 \pm 7.2 \\ -1 \pm 12 \\ -2 \pm 21 \\ 3.0 \pm 2.3 \end{pmatrix} \times 10^{-2}$	$\begin{pmatrix} [c_{\ell q}^{(3)}]_{1111} \\ [c_{eq}]_{1111} \\ [\hat{c}_{\ell u}]_{1111} \\ [\hat{c}_{\ell d}]_{1111} \\ [c_{eu}]_{1111} \\ [\hat{c}_{ed}]_{1111} \\ [\hat{c}_{\ell q}^{(3)}]_{1122} \\ [c_{\ell u}]_{1122} \\ [\hat{c}_{\ell d}]_{1122} \\ [c_{eq}]_{1122} \\ [c_{eu}]_{1122} \\ [c_{ed}]_{1122} \\ [\hat{c}_{\ell q}^{(3)}]_{1133} \\ [c_{\ell u}]_{1133} \\ [c_{eq}]_{1133} \\ [c_{eu}]_{1133} \\ [c_{\ell q}^{(3)}]_{2211} \\ [c_{\ell u}]_{2211} \\ [c_{\ell d}]_{2211} \\ [c_{eq}]_{2211} \\ [c_{\ell qu}]_{1111} \\ [c_{\ell dq}]_{1111} \\ [c_{\ell qu}^{(3)}]_{1111} \\ c_P^{d\mu} (2 \text{ GeV}) \end{pmatrix} = \begin{pmatrix} -2.2 \pm 3.2 \\ 100 \pm 180 \\ -5 \pm 11 \\ -5 \pm 23 \\ -1 \pm 12 \\ -4 \pm 21 \\ -61 \pm 32 \\ 2.4 \pm 8.0 \\ -310 \pm 130 \\ -21 \pm 28 \\ -87 \pm 46 \\ 270 \pm 140 \\ -8.6 \pm 8.0 \\ -1.4 \pm 10 \\ -3.2 \pm 5.1 \\ 18 \pm 20 \\ -1.2 \pm 3.9 \\ 1.3 \pm 7.6 \\ 15 \pm 12 \\ 25 \pm 34 \\ 4 \pm 41 \\ -0.080 \pm 0.075 \\ -0.079 \pm 0.074 \\ -0.02 \pm 0.19 \\ -0.02 \pm 0.15 \end{pmatrix} \times 10^{-2}$	 
--	---	--	--