

How to (try? hope? to) resolve the Hubble tension

Vivian Poulin

Laboratoire Univers et Particules de Montpellier
CNRS & Université de Montpellier


w/ T. Smith, T. Karwal, M. Kamionkowski, M. Amin
A. Klypin, F. Prada, J. Primack, V. Avila-Reese, A. Rodriguez-Puebla, P. Behroozi, D. Hellinger
R. Murgia, G. F. Abellan

Instituto de Física Corpuscular, Valencia
March, 9th 2021



The Era of Precision Cosmology


20 years ago



© Dodelson


The Era of Precision Cosmology

10 years ago




The Era of Precision Cosmology

Today




The Era of Precision Cosmology

Good agreement between all CMB data!



The Era of Precision Cosmology

And also with non-CMB data!



The Era of Precision Cosmology

Astonishing success of Λ CDM Cosmology

	Parameter	<i>Planck</i> alone	<i>Planck</i> + BAO	<u>Planck alone</u>
matter content	$\Omega_b h^2$	0.02237 ± 0.00015	0.02242 ± 0.00014	0.6% precision
expansion rate (H_0, Λ)	$\Omega_c h^2$	0.1200 ± 0.0012	0.11933 ± 0.00091	1% precision
	$100\theta_{MC}$	1.04092 ± 0.00031	1.04101 ± 0.00029	0.3% precision
star formation	τ	0.0544 ± 0.0073	0.0561 ± 0.0071	13% precision
	$\ln(10^{10} A_s)$	3.044 ± 0.014	3.047 ± 0.014	5% precision
Inflation	n_s	0.9649 ± 0.0042	0.9665 ± 0.0038	0.5% precision
	H_0	67.36 ± 0.54	67.66 ± 0.42	0.7% precision


e.g. 2015 data: TT +lowP reduced $\chi^2 = 1.004$

As precision of data has increased, a certain number of “tensions” have emerged

- Hubble tension: Direct H_0 measurements are higher than Λ CDM prediction.
- $S_8 = \sigma_8(\Omega_m/0.3)^{0.5}$ prediction is higher at $\sim 2\text{-}3\sigma$ than that measured by Weak Lensing surveys (CFHTLenS, KiDS, DES, HSC).
- Internal tension(s) within Planck and with other CMB datasets $\sim 2\text{-}3\sigma$.

See ‘Cosmology intertwined’ Di Valentino, Anchordoqui++ 2008.1128(3-6), Handley&Lemos 2007.08496

The Hubble tension




In 2017: $\sim 3.5\sigma$ discrepancy between "direct" measurement from SH0ES and the value inferred from a fit of ΛCDM to *Planck* 2015


The Hubble tension

Since then, over 20 measurements
and 800 papers!!

- Indirect: H_0 is a prediction from the Λ CDM model constrained with high-z data
- Direct: H_0 is measured at low-z in different ways
- All Direct measurements are higher than Indirect ones, not all are in strong tension
- Average: tension between $4\text{--}6\sigma$



The Hubble Constant in 3 Steps: Present Data



Systematics in the ‘SN1a’ route?


- Tension: -0.2μ shift is on the order of the scatter σ *Freedman et al., Apj 2019*

- Discussion about the amount of dust extinction towards the LMC between CCHP and SH0ES team.


Freedman et al., 2019. Yuan et al 2019, Freedman et al. 2020

- Latest calibration of TRGB with GAIA DR3:
 $H_0 = 72.1 \pm 2 \text{ km/s/Mpc}$

Soltis et al 2020




TRGB – SNIa	
Soltis, Casertano, Riess (2020):	72.1 ± 2.0
Freedman et al. (2020):	69.6 ± 1.9
Reid, Pesce, Riess (2019), SH0ES:	71.1 ± 1.9
Freedman et al. (2019):	69.8 ± 1.9
Yuan et al. (2019):	72.4 ± 2.0
Jang, Lee (2017):	71.2 ± 2.5




H0LiCOW: QSOs gravitational time delay


$$D_{\Delta t} \equiv (1 + z_d) \frac{D_d D_s}{D_{ds}}$$




(a) B1608+656




(b) RXJ1131-1231




(c) HE 0435-1223



(d) SDSS 1206+4332




(e) WFI2033-4723




(f) PG 1115+080


Wong et al. 1907.04869



H0LiCOW: H_0 measurement to few %



- 6 QSOs: 3.1σ tension with *Planck* within Λ CDM
- Blind analysis
- This assumes a specific shape for the lens mass profile.





Wong et al. 1907.04869

TDCOSMO+SLAC: Agreement with Λ CDM?


Birrer++ 2007.02941

H_0 measurements in flat Λ CDM - performed blindly



- New analysis relaxes the power-law profile assumption and rely on stellar kinematics.
- Same sample of galaxies (TDCOSMO): large H_0 but large error bars.
- Add information from the ‘SLACS’ lenses catalog: 5% error and H_0 decreases.

H_0 : the pessimistic (realistic?) view



© Colin Hill

- The situation is somewhat complicated... Hopefully resolved in ~year time scale.

How does CMB data measure H_0 ?

- The ‘sound horizon’ r_s , a standard ruler in the sky: distance travelled by sound wave until recombination.
- Planck measures θ_s and, given a model, can extract r_s .
- H_0 appears *only* in the angular diameter distance d_A .

$$\theta_s \equiv \frac{r_s(z_*)}{d_A(z_*)} = \frac{\int_{\infty}^{z_*} dz \, c_s(z) / \sqrt{\rho_{\text{tot}}(z)}}{\int_0^{z_*} dz / \sqrt{\rho_{\text{tot}}(z)}}$$




illustration: T. Smith


Geometrical degeneracy in Planck!

- A higher H_0 can be compensated by a lower $H(z > 0)$ such as to keep $d_A(z_*)$ fixed

$$d_A(z_*) = \frac{1}{1+z_*} \int_0^{z_*} \frac{dz}{100 \sqrt{\omega_M(1+z)^3 + \Omega_{DE}(z)h^2}}$$

- ‘phantom dark energy’ $w < -1$, DE phase transition, DE-DM interaction, decaying/annihilating DM, and many more...
[\[http://arxiv.org/insert_your_favorite_model_here.com\]](http://arxiv.org/insert_your_favorite_model_here.com)

- Planck can easily accommodate a higher H_0 : problem with BAO and Pantheon






BAO and SN1a constrain late-time resolution

$$d_A(z_*) = \frac{1}{1+z_*} \int_0^{z_*} \frac{dz}{100 \sqrt{\omega_M(1+z)^3 + \Omega_{DE}(z)h^2}}$$

VP, Boddy, Bird, Kamionkowski 1803.02474

see also Wang++ 1807.03772, Bernal++ 1607.05617,
Raveri 1902.01366, Aylor++1811.00537, Benevento++ 2002.11707.



Knox&Millea 1908.03663

$$\theta_d(z)^\perp = \frac{r_s(z_{\text{drag}})}{D_A(z)}, \theta_d(z)^\parallel = r_s(z_{\text{drag}})H(z)$$

- $r_s(z_{\text{drag}})$ from Planck

$$\mu(z) = 5\text{Log}_{10}D_L(z) + \text{const.}$$

- Calibration constant from e.g. SH0ES.


In GR: $D_A = D_L/(1+z)^2$; it is impossible to resolve the tension without changing calibration!

- Nb: unless of a ‘cosmological crisis’ *Di Valentino++ 1911.02087*

H_0 tension or r_s tension?

One can deduce the co-moving sound horizon r_s from H_0 and BAO

r_s from CMB needs to **decrease by ~ 10 Mpc**



A sketch of the physics at play

- Could the CMB be closer to us than Λ CDM tells us? This is what a higher H_0 suggests.
- Therefore, could spot in the CMB be smaller? This is what new physics must achieve.




illustration: T. Smith

Early-time resolution to the H_0 tension

affect z_* : modified recombination physics?

Chiang&Slosar 1811.03624

see also *Hart&Chluba 1912.03986*

Jedamzik&Pogosian 2004.09487


affect c_s : DM-photon scattering? DM-b scattering?

Boddy, Gluscevic, VP++1808.00001


$$r_s = \int_{\infty}^{z_*} dz \frac{c_s(z)}{H(z)}$$

affect $H(z)$: N_{eff} ? Early Dark Energy?

- N_{eff} (free streaming) $\sim 3.5-4$ is needed : disfavored by *Planck* high- l polarization and BAO



Bernal++ 1607.05617




Aghanim++ 1807.06209

- Interacting neutrinos might work, although TBC with Planck 18 *Kreisch++ 1902.00534*

Early-Universe solution to H_0

- r_s does not reach **10Mpc** before ~ 25000 in Λ CDM



GOAL: decreasing r_s by 10Mpc while keeping r_s/r_d and r_s/r_{eq} fixed

Early Dark Energy(s) is a promising solution

Early dark energy, the Hubble-parameter tension, and the string axiverse

Tanvi Karwal and Marc Kamionkowski
Department of Physics and Astronomy, Johns Hopkins University,
3400 N. Charles St., Baltimore, MD 21218
(Dated: November 8, 2016)

Not all have the same
success...

Early Dark Energy Can Resolve The Hubble Tension

Vivian Poulin¹, Tristan L. Smith², Tanvi Karwal¹, and Marc Kamionkowski¹
¹Department of Physics and Astronomy, Johns Hopkins University,
3400 N. Charles St., Baltimore, MD 21218, United States and
²Department of Physics and Astronomy, Swarthmore College,
500 College Ave., Swarthmore, PA 19081, United States

Rock 'n' Roll Solutions to the Hubble Tension

Prateek Agrawal¹, Francis-Yan Cyr-Racine^{1,2}, David Pinner^{1,3}, and Lisa Randall¹

¹Department of Physics, Harvard University, 17 Oxford St., Cambridge, MA 02138, USA

²Department of Physics and Astronomy, University of New Mexico, 1919 Lomas Blvd NE, Albuquerque, NM 87131, USA

³Department of Physics, Brown University, 182 Hope St., Providence, RI 02912, USA

Acoustic Dark Energy: Potential Conversion of the Hubble Tension

Meng-Xiang Lin,¹ Giampaolo Benevento,^{2,3,1} Wayne Hu,¹ and Marco Raveri¹

¹Kavli Institute for Cosmological Physics, Department of Astronomy & Astrophysics,
Enrico Fermi Institute, The University of Chicago, Chicago, IL 60637, USA

²Dipartimento di Fisica e Astronomia "G. Galilei",

Università degli Studi di Padova, via Marzolo 8, I-35131, Padova, Italy

³INFN, Sezione di Padova, via Marzolo 8, I-35131, Padova, Italy

Early dark energy from massive neutrinos — a natural resolution of the Hubble tension

Jeremy Sakstein* and Mark Trodden[†]

Center for Particle Cosmology, Department of Physics and Astronomy,
University of Pennsylvania 209 S. 33rd St., Philadelphia, PA 19104, USA

Dark Energy, H_0 and Weak Gravity Conjecture

Nemanja Kaloper^{a,1}

^aDepartment of Physics, University of California, Davis, CA 95616, USA

Is the Hubble tension a hint of AdS around recombination?

Gen Ye^{1*} and Yun-Song Piao^{1,2†}

¹ School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China and

² Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100190, China

Thermal Friction as a Solution to the Hubble Tension

Kim V. Berghaus¹ and Tanvi Karwal^{1,2}

¹Department of Physics and Astronomy, Johns Hopkins University,
3400 N. Charles St., Baltimore, MD 21218, United States and

²Center for Particle Cosmology, Department of Physics and Astronomy,
University of Pennsylvania, 209 S. 33rd St., Philadelphia, PA 19104, United States

(Dated: November 15, 2019)

Scalar-tensor theories of gravity, neutrino physics, and the H_0 tension

Mario Ballardini,^{a,b,c,d,1} Matteo Braglia,^{a,b,c} Fabio Finelli,^{b,c} Daniela Paoletti,^{b,c} Alexei A. Starobinsky,^{e,f} Caterina Umiltà^g

New Early Dark Energy

Florian Niedermann^{1,*} and Martin S. Sloth^{1,†}

CP³-Origins, Center for Cosmology and Particle Physics Phenomenology

Gravity in the Era of Equality: Towards solutions to the Hubble problem without fine-tuned initial conditions

Miguel Zumalacárregui^{1,2,3,*}

¹Max Planck Institute for Gravitational Physics (Albert Einstein Institute)
Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany

²Berkeley Center for Cosmological Physics, LBNL and University of California at Berkeley,
Berkeley, California 94720, USA

³Institut de Physique Théorique, Université Paris Saclay CEA, CNRS, 91191 Gif-sur-Yvette, France
(Dated: June 11, 2020)

Scalar field and Early Dark Energy

- Initially **slowly-rolling field** (due to Hubble friction) that later **dilutes faster than matter**

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV_n(\phi)}{d\phi} = 0 \quad \rho_\phi = \frac{1}{2}\dot{\phi}^2 + V_n(\phi), \quad P_\phi = \frac{1}{2}\dot{\phi}^2 - V_n(\phi)$$

- Oscillating (toy) potential:


$$V(\phi) \propto (1 - \cos \phi)^n$$

VP++ 1806.10608 & 1811.04083
Smith, VP++ 1908.06995

- Specified by $f_{\text{EDE}}(z_c)$, z_c , $w(n)$, $c_s^2(k, \tau)$

$$\begin{cases} z > z_c \Rightarrow w_n = 1 \\ z < z_c \Rightarrow w_n = (n - 1)/(n + 1) \end{cases}$$


$n = 1$: matter, $n = 2$: radiation, etc.



See also Niedermann&Sloth for different EDE model 1910.10739, 2006.06686, 2009.00006

EDE Can Resolve The Hubble Tension

VP++PRL 122 (2019); Smith++1908.06995



- Planck high- ℓ TT,TE, EE+lowTEB+lensing+BAO+Pantheon+SH0ES 19


$$f(z_c) = 0.10 \text{ (0.13)} \pm 0.03 \quad \text{Log}_{10}(z_c) = 3.56 \text{ (3.53)}^{+0.05}_{-0.1} \quad H_0 = 71.5 \text{ (72.8)} \pm 1.2 \text{ km/s/Mpc}$$

- $n < 3.5$ at 1σ : scalar field **oscillations are favored** over non-oscillating solutions

Datasets	Λ CDM	n free
Planck high- ℓ TT, TE, EE	2446.66	2445.53
Planck low- ℓ TT, TE, EE	10496.65	10493.65
Planck lensing	10.37	9.14
SH0ES	16.80	0.73
Total $\chi^2_{\min}$	14001.23	13980.90
$\Delta\chi^2_{\min}$	0	-20.33


See also recent update in Murgia, Abellan, VP 2009.10733

EDE leaves an imprint in CMB power spectra



A New Understanding Of Λ ?

- The field becomes dynamical around z_{eq} : Fine tuning ? Coincidence problem 2.0?
- What if there were **more of such era to be discovered**? We already have seen two (three?) of them.
- Is there one field with a complicated potential or many fields with simple potentials?
e.g. Dodelson++astro-ph/0002360, Griest astro-ph/0202052, Kamionkowski++1409.0549




CMB data can constrain c_s^2

- In Lin, Beneveto, Hu, Raveri 1905.12618: “Acoustic” Dark Energy with constant c_s^2 .

$$P(X, \phi) = \left(\frac{X}{A}\right)^{\frac{1-c_s^2}{2c_s^2}} X - V(\phi),$$

$$V(\phi) = \begin{cases} A\phi^m, & \phi > 0, \\ 0, & \phi \leq 0. \end{cases}$$



For $w = 0.5$ ($n=3$) data favors $c_s^2 < 0.9$ at 95% C.L.

Detecting the EDE with CMB data only


- Future CMB experiment like CMB-S4 **will be able to detect the EDE** without SH0ES data.

Fiducial model:


$$f(z_c) = 0.12$$

$$z_c = 10^{3.5}$$

$$h = 0.72$$





- Without including the EDE: one might **strongly bias** H_0 and ω_{cdm} values.



$$S_8 \equiv \sigma_8(\Omega_m/0.3)^{0.5}$$



The S_8 tension



- $\sim 2.5\sigma$ tension between **CFHTLenS/HSC/KiDS** and *Planck*. (Also Planck SZ?)
Heymans++ MNRAS 2013, KiDS MNRAS 2017, HSC PASJ 2019, DES PRD 2018, Salvati++ PoS 1901.05289
- **DES galaxies or shear alone in agreement** with *Planck*, while **galaxies+shear** shows 2σ tension. Hot discussion about calibration of photometric redshift.
DES collab. PRD 2018, Joudaki++ 1906.09262
- To resolve the tension: reduce power at scales $k \gtrsim 0.1$ h/Mpc. DM interactions, decay, massive neutrinos, (warm or fuzzy dark matter?)
[insert your favorite paper here]

EDE can be constrained by LSS



Murgia, Abellan, VP 2009.10733



- The increase in ω_{cdm} & n_s leads to an increase in S_8 .
- The S_8 tension increases by $\sim 0.5\sigma$.
- Constraints from KIDS? DES? BOSS?

EDE can be constrained by LSS



Murgia, Abellan, VP 2009.10733



- Including BOSS RSD measurements of $f\sigma_8$:
 $f_{\text{EDE}}(a_c) = 0.097^{+0.035}_{-0.029}$
 $H_0 = 71^{+1.1}_{-1}$
- $\Delta\chi^2 \simeq -20$ (stable)

EDE can be constrained by LSS


Murgia, Abellan, VP 2009.10733



- Combining with KiDS/KV(+H0):
 $f_{\text{EDE}} = 0.085^{+0.034}_{-0.022}$ $H_0 = 70.87^{+1}_{-0.91}$
- KIDS/VIKING/DES(-H0):
 $f_{\text{EDE}}(z_c) < 0.06$ $H_0 = 68.92^{+0.57}_{-0.59}$
Hill et al. 2003.07355
- Statistical inconsistency!
 'compromise' cosmology is a bad fit to S_8 data *Raveri&Hu 1806.04649*

Could the A_L anomaly be related to S_8 ?

Murgia, Abellan, VP 2009.10733





- Marginalize over lensing information in *Planck*: Lensing power spectrum amplitude A_L + peak smoothing amplitude A_L^{TTTEEE}
- ‘ A_L anomaly’: The lensing amplitude deduced from peak smoothing A_L^{TTTEEE} in *Planck* data is higher than that from lensing reconstruction A_L .
- In Λ CDM: extra smoothing requires higher A_s and $\omega_{\text{cdm}} \Rightarrow$ higher S_8 .
- Once marginalizing over the ‘ A_L anomaly’; no S_8 tension! no anomaly in ACT&SPT.

see also Di Valentino & Bridle 2018, *Planck* 1807.06209, *SPTPol* 1910.07157


Lensing-marginalized EDE cosmology vs S_8

Murgia, Abellan, VP 2009.10733





- Once marginalizing over the A_L anomaly; EDE in agreement with S_8 data!
- The determination of A_L and A_L^{TTTEEE} is unaffected by the EDE parameters!
- This is also supported by SPTPol analysis.

No S8-tension with ACT and SPT



What will it take to find a concordance model?

Jedamzik & Pogosian 2010.04158



- Fitting Planck, H_0 and BAO with lower $r_s(z_{\text{rec}})$ requires higher ω_{cdm} : the S_8 tension increases.
- Resolving both tensions (unless systematics!) will likely require multiple extensions
- H_0 : measure the background expansion rate. S_8 : measure the amplitude of perturbations.
- It is likely that a solution will need some specific background & perturbation dynamics.
- This is already the case in a variety of model! Interacting DM-DR, interacting neutrinos, EDE- m_ν model... But very limited success so-far.

Conclusions

- H_0 from local measurements is in tension with LCDM-inferred value from Planck. Systematics are still possible!
- This is better understood when recast as a **sound-horizon tension**: CMB r_s too high by 10Mpc.
- A frozen scalar field acting like **Early Dark Energy until $z_c \sim 3500$ with $f(z_c) \sim 10\%$** and diluting faster than radiation later can solve the Hubble tension.
- Slightly better fit to Planck data, once SH0ES is included **“definite” evidence for $n \geq 3$, $\Delta\chi^2_{\min} \sim -20$.**
- Future CMB data will unambiguously detect (or exclude) EDE. If this is the “correct” resolution: there might be **new ways of interpreting Λ** and inflation.
- This naive **model is disfavored (not ruled out!)** by recent KiDS and BOSS data. Combining Planck and LSS data might not be statistically sound! S_8 tension, A_L anomalous, A_s discrepancy.
- Connections between S_8 and H_0 exist but are not automatic (perturbations vs background).

Thanks for your attention!