

A_{FB} in the SMEFT: the LHC as a Z physics laboratory

Víctor Bresó-Pla
IFIC, CSIC/U. Valencia

In collaboration with

Adam Falkowski and Martín González-Alonso

arXiv:[2103.12074]



VNIVERSITAT
D VALÈNCIA



CSIC
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

1. Introduction


The indirect search for Beyond Standard Model (BSM) physics requires very precise measurements and predictions and greatly benefits from the use of a general framework

1. Introduction

The indirect search for Beyond Standard Model (BSM) physics requires very precise measurements and predictions and greatly benefits from the use of a general framework



General framework: SMEFT



1. Introduction

The indirect search for Beyond Standard Model (BSM) physics requires very precise measurements and predictions and greatly benefits from the use of a general framework



Precise measurements: LEP + LHC

- LEP provides very precise measurements which leave little room for NP at the EW scale. LHC and Tevatron have also contributed to this thanks to some of their measurements for observables in the W, top and Higgs sectors

1. Introduction

The indirect search for Beyond Standard Model (BSM) physics requires very precise measurements and predictions and greatly benefits from the use of a general framework



Precise measurements: LEP + LHC

- LEP provides very precise measurements which leave little room for NP at the EW scale. LHC and Tevatron have also contributed to this thanks to some of their measurements for observables in the W, top and Higgs sectors
- **Could it be that LHC can also compete with or complement LEP by means of the measurements it can provide for the Z observables?**

1. Introduction

The indirect search for Beyond Standard Model (BSM) physics requires very precise measurements and predictions and greatly benefits from the use of a general framework



Precise measurements: LEP + LHC

- LEP provides very precise measurements which leave little room for NP at the EW scale. LHC and Tevatron have also contributed to this thanks to some of their measurements for observables in the W, top and Higgs sectors
- **Could it be that LHC can also compete with or complement LEP by means of the measurements it can provide for the Z observables?**
- We will explore this possibility by looking at Drell-Yan dilepton production, which are sensitive to the Zff couplings

2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\begin{aligned}
 \mathcal{L}_{\text{SMEFT}} \supset & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\
 & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right) \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right] \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]
 \end{aligned}$$

$$\mathcal{L}_{\text{SMEFT}} \supset \frac{g_L^2 v^2}{4} (1 + \delta m_w)^2 W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{8} Z_\mu Z_\mu$$

2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\begin{aligned}
 \mathcal{L}_{\text{SMEFT}} \supset & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\
 & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right) \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right] \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]
 \end{aligned}$$

Input scheme:
 $\{G_F, \alpha(M_Z), M_Z\}$

$$\mathcal{L}_{\text{SMEFT}} \supset \frac{g_L^2 v^2}{4} (1 + \delta m_w)^2 W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{8} Z_\mu Z_\mu$$

2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\begin{aligned}
 \mathcal{L}_{\text{SMEFT}} \supset & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\
 & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right) \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right] \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]
 \end{aligned}$$

Input scheme:
 $\{G_F, \alpha(M_Z), M_Z\}$

$$\mathcal{L}_{\text{SMEFT}} \supset \frac{g_L^2 v^2}{4} (1 + \delta m_w)^2 W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{8} Z_\mu Z_\mu$$

Flavor-general corrections!!

[Efrati, Falkowski & Soreq '15;
 Falkowski & Mimouni '15; Falkowski,
 González-Alonso & Mimouni '17;
 Aebischer et al. '18; ...]


2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\mathcal{L}_{\text{SMEFT}} \supset -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\ -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right)$$

$$-\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right]$$

$$\left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]$$



$$(\delta m_w)^2 W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{8} Z_\mu Z_\mu$$

Flavor-general corrections!!

[Efrati, Falkowski & Soreq '15;
Falkowski & Mimouni '15; Falkowski,
González-Alonso & Mimouni '17;
Aebischer et al. '18; ...]

2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\begin{aligned}
 \mathcal{L}_{\text{SMEFT}} \supset & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\
 & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right) \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right] \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]
 \end{aligned}$$

$$\mathcal{L}_{\text{SMEFT}} \supset \frac{g_L^2 v^2}{4} (1 + \delta m_w)^2 W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{8} Z_\mu Z_\mu$$

$$\delta g_L^{We}, \delta g_L^{W\mu}, \delta g_L^{W\tau}, \delta g_{L/R}^{Ze}, \delta g_{L/R}^{Z\mu}, \delta g_{L/R}^{Z\tau}, \delta g_{L/R}^{Zd}, \delta g_{L/R}^{Zs}, \delta g_{L/R}^{Zb}, \delta g_{L/R}^{Zu}, \delta g_{L/R}^{Zc}, \delta m_w$$

2. Theory framework

- **SMEFT**: We focus only on **Z and W pole observables**, which are mainly sensitive to non-derivative interactions between EW bosons and fermions:

$$\begin{aligned}
 \mathcal{L}_{\text{SMEFT}} \supset & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{u}_L \gamma_\mu (V + \delta g_L^{Wq}) d_L + W_\mu^+ \bar{u}_R \gamma_\mu \delta g_R^{Wq} d_R + \text{h.c.} \right) \\
 & -\frac{g_L}{\sqrt{2}} \left(W_\mu^+ \bar{\nu}_L \gamma_\mu (\mathbf{I} + \delta g_L^{We}) e_L + \text{h.c.} \right) \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e, \nu} \bar{f}_L \gamma_\mu ((T_f^3 - s_\theta^2 Q_f) \mathbf{I} + \delta g_L^{Zf}) f_L \right] \\
 & -\sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u, d, e} \bar{f}_R \gamma_\mu (-s_\theta^2 Q_f \mathbf{I} + \delta g_R^{Zf}) f_R \right]
 \end{aligned}$$

$$\mathcal{L}_{\text{SMEFT}} \supset \frac{g_L^2 v^2}{4} (1 + \delta m_w)^2 W_\mu^+ W_\mu^- + \dots$$

We end up with only 20 independent parameters

$$\delta g_L^{We}, \delta g_L^{W\mu}, \delta g_L^{W\tau}, \delta g_{L/R}^{Ze}, \delta g_{L/R}^{Z\mu}, \delta g_{L/R}^{Z\tau}, \delta g_{L/R}^{Zd}, \delta g_{L/R}^{Zs}, \delta g_{L/R}^{Zb}, \delta g_{L/R}^{Zu}, \delta g_{L/R}^{Zc}, \delta m_w$$

3. “Traditional” pole observables

○ Z pole observables:

Observable	Experimental value	SM prediction	Definition
Γ_Z [GeV]	2.4955 ± 0.0023 [4, 28]	2.4941	$\sum_f \Gamma(Z \rightarrow f\bar{f})$
σ_{had} [nb]	41.4802 ± 0.0325 [4, 28]	41.4842	$\frac{12\pi}{m_Z^2} \frac{\Gamma(Z \rightarrow e^+e^-)\Gamma(Z \rightarrow q\bar{q})}{\Gamma_Z^2}$
R_e	20.804 ± 0.050 [4]	20.734	$\frac{\sum_q \Gamma(Z \rightarrow q\bar{q})}{\Gamma(Z \rightarrow e^+e^-)}$
R_μ	20.785 ± 0.033 [4]	20.734	$\frac{\sum_q \Gamma(Z \rightarrow q\bar{q})}{\Gamma(Z \rightarrow \mu^+\mu^-)}$
R_τ	20.764 ± 0.045 [4]	20.781	$\frac{\sum_q \Gamma(Z \rightarrow q\bar{q})}{\Gamma(Z \rightarrow \tau^+\tau^-)}$
$A_{\text{FB}}^{0,e}$	0.0145 ± 0.0025 [4]	0.0162	$\frac{3}{4}A_e^2$
$A_{\text{FB}}^{0,\mu}$	0.0169 ± 0.0013 [4]	0.0162	$\frac{3}{4}A_e A_\mu$
$A_{\text{FB}}^{0,\tau}$	0.0188 ± 0.0017 [4]	0.0162	$\frac{3}{4}A_e A_\tau$
R_b	0.21629 ± 0.00066 [4]	0.21581	$\frac{\Gamma(Z \rightarrow b\bar{b})}{\sum_q \Gamma(Z \rightarrow q\bar{q})}$
R_c	0.1721 ± 0.0030 [4]	0.17222	$\frac{\Gamma(Z \rightarrow c\bar{c})}{\sum_q \Gamma(Z \rightarrow q\bar{q})}$
A_b^{FB}	0.0996 ± 0.0016 [4, 29]	0.1032	$\frac{3}{4}A_e A_b$
A_c^{FB}	0.0707 ± 0.0035 [4]	0.0736	$\frac{3}{4}A_e A_c$
A_e	0.1516 ± 0.0021 [4]	0.1470	$\frac{\Gamma(Z \rightarrow e_L^+e_L^-) - \Gamma(Z \rightarrow e_R^+e_R^-)}{\Gamma(Z \rightarrow e^+e^-)}$
A_μ	0.142 ± 0.015 [4]	0.1470	$\frac{\Gamma(Z \rightarrow \mu_L^+\mu_L^-) - \Gamma(Z \rightarrow \mu_R^+\mu_R^-)}{\Gamma(Z \rightarrow \mu^+\mu^-)}$
A_τ	0.136 ± 0.015 [4]	0.1470	$\frac{\Gamma(Z \rightarrow \tau_L^+\tau_L^-) - \Gamma(Z \rightarrow \tau_R^+\tau_R^-)}{\Gamma(Z \rightarrow \tau^+\tau^-)}$
A_e	0.1498 ± 0.0049 [4]	0.1470	$\frac{\Gamma(Z \rightarrow e_L^+e_L^-) - \Gamma(Z \rightarrow e_R^+e_R^-)}{\Gamma(Z \rightarrow e^+e^-)}$
A_τ	0.1439 ± 0.0043 [4]	0.1470	$\frac{\Gamma(Z \rightarrow \tau_L^+\tau_L^-) - \Gamma(Z \rightarrow \tau_R^+\tau_R^-)}{\Gamma(Z \rightarrow \tau^+\tau^-)}$
A_b	0.923 ± 0.020 [4]	0.935	$\frac{\Gamma(Z \rightarrow b_L b_L) - \Gamma(Z \rightarrow b_R b_R)}{\Gamma(Z \rightarrow b\bar{b})}$
A_c	0.670 ± 0.027 [4]	0.668	$\frac{\Gamma(Z \rightarrow c_L c_L) - \Gamma(Z \rightarrow c_R c_R)}{\Gamma(Z \rightarrow c\bar{c})}$
A_s	0.895 ± 0.091 [30]	0.936	$\frac{\Gamma(Z \rightarrow s_L s_L) - \Gamma(Z \rightarrow s_R s_R)}{\Gamma(Z \rightarrow s\bar{s})}$
R_{uc}	0.166 ± 0.009 [9]	0.1722	$\frac{\Gamma(Z \rightarrow u\bar{u}) + \Gamma(Z \rightarrow c\bar{c})}{2 \sum_q \Gamma(Z \rightarrow q\bar{q})}$

○ W pole observables:

Observable	Experimental value	SM prediction
m_W [GeV]	80.379 ± 0.012 [9]	80.356
Γ_W [GeV]	2.085 ± 0.042 [9]	2.088
$\text{Br}(W \rightarrow e\nu)$	0.1071 ± 0.0016 [5]	0.1082
$\text{Br}(W \rightarrow \mu\nu)$	0.1063 ± 0.0015 [5]	0.1082
$\text{Br}(W \rightarrow \tau\nu)$	0.1138 ± 0.0021 [5]	0.1081
$\text{Br}(W \rightarrow \mu\nu)/\text{Br}(W \rightarrow e\nu)$	0.982 ± 0.024 [32]	1.000
$\text{Br}(W \rightarrow \mu\nu)/\text{Br}(W \rightarrow e\nu)$	1.020 ± 0.019 [12]	1.000
$\text{Br}(W \rightarrow \mu\nu)/\text{Br}(W \rightarrow e\nu)$	1.003 ± 0.010 [13]	1.000
$\text{Br}(W \rightarrow \tau\nu)/\text{Br}(W \rightarrow e\nu)$	0.961 ± 0.061 [9, 31]	0.999
$\text{Br}(W \rightarrow \tau\nu)/\text{Br}(W \rightarrow \mu\nu)$	0.992 ± 0.013 [14]	0.999
$R_{Wc} \equiv \frac{\Gamma(W \rightarrow cs)}{\Gamma(W \rightarrow ud) + \Gamma(W \rightarrow cs)}$	0.49 ± 0.04 [9]	0.50

3. “Traditional” pole observables

- Leptonic couplings:

$$[\delta g_L^{We}]_{ii} = \begin{pmatrix} -1.3 \pm 3.2 \\ -2.8 \pm 2.6 \\ 1.5 \pm 4.0 \end{pmatrix} \times 10^{-3}$$

$$[\delta g_R^{Ze}]_{ii} = \begin{pmatrix} -0.43 \pm 0.27 \\ 0.0 \pm 1.4 \\ 0.62 \pm 0.62 \end{pmatrix} \times 10^{-3}$$

$$[\delta g_L^{Ze}]_{ii} = \begin{pmatrix} -0.19 \pm 0.28 \\ 0.1 \pm 1.2 \\ -0.09 \pm 0.59 \end{pmatrix} \times 10^{-3}$$

- W mass correction:

$$\delta m_w = (2.9 \pm 1.6) \times 10^{-4}$$

- s, c, b couplings:

$$\delta g_L^{Zs} = (1.3 \pm 4.1) \times 10^{-2}$$

$$\delta g_R^{Zs} = (2.2 \pm 5.6) \times 10^{-2}$$

$$\delta g_L^{Zc} = (-1.3 \pm 3.7) \times 10^{-3}$$

$$\delta g_R^{Zc} = (-3.2 \pm 5.4) \times 10^{-3}$$

$$\delta g_L^{Zb} = (3.1 \pm 1.7) \times 10^{-3}$$

$$\delta g_R^{Zb} = (21.8 \pm 8.8) \times 10^{-3}$$

[Update of Efrati, Falkowski & Soreq '15]

3. “Traditional” pole observables

- Leptonic couplings:

$$[\delta g_L^{We}]_{ii} = \begin{pmatrix} -1.3 \pm 3.2 \\ -2.8 \pm 2.6 \\ 1.5 \pm 4.0 \end{pmatrix} \times 10^{-3}$$

$$[\delta g_R^{Ze}]_{ii} = \begin{pmatrix} -0.43 \pm 0.27 \\ 0.0 \pm 1.4 \\ 0.62 \pm 0.62 \end{pmatrix} \times 10^{-3}$$

$$[\delta g_L^{Ze}]_{ii} = \begin{pmatrix} -0.19 \pm 0.28 \\ 0.1 \pm 1.2 \\ -0.09 \pm 0.59 \end{pmatrix} \times 10^{-3}$$

- W mass correction:

$$\delta m_w = (2.9 \pm 1.6) \times 10^{-4}$$

- s, c, b couplings:

What about Zuu and Zdd corrections?

$$\delta g_L^{Zs} = (1.3 \pm 4.1) \times 10^{-2}$$

$$\delta g_R^{Zs} = (2.2 \pm 5.6) \times 10^{-2}$$

$$\delta g_L^{Zc} = (-1.3 \pm 3.7) \times 10^{-3}$$

$$\delta g_R^{Zc} = (-3.2 \pm 5.4) \times 10^{-3}$$

$$\delta g_L^{Zb} = (3.1 \pm 1.7) \times 10^{-3}$$

$$\delta g_R^{Zb} = (21.8 \pm 8.8) \times 10^{-3}$$

[Update of Efrati, Falkowski & Soreq '15]

3. “Traditional” pole observables

- One linear combination of up and down quark vertex corrections is unconstrained:

$$\delta g_L^{Zu} + \delta g_L^{Zd} + \frac{3g_L^2 - g_Y^2}{4g_Y^2} \delta g_R^{Zu} + \frac{3g_L^2 + g_Y^2}{2g_Y^2} \delta g_R^{Zd}$$

- It is useful to rearrange these 4 couplings so that we can separate the blind direction from the rest of the parameter space:

$$\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = R \begin{pmatrix} \delta g_L^{Zu} \\ \delta g_R^{Zu} \\ \delta g_L^{Zd} \\ \delta g_R^{Zd} \end{pmatrix} = \begin{pmatrix} 0.93 & -0.29 & -0.23 & -0.01 \\ 0.18 & 0.87 & -0.33 & -0.33 \\ 0.27 & 0.18 & 0.90 & -0.29 \\ 0.17 & 0.37 & 0.17 & 0.90 \end{pmatrix} \begin{pmatrix} \delta g_L^{Zu} \\ \delta g_R^{Zu} \\ \delta g_L^{Zd} \\ \delta g_R^{Zd} \end{pmatrix}$$



$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.9 \pm 1.8 \\ 0.3 \pm 3.3 \\ -2.4 \pm 4.8 \end{pmatrix} \times 10^{-2}$$

3. “Traditional” pole observables

- One linear combination of up and down quark vertex corrections is unconstrained:

$$\delta g_L^{Zu} + \delta g_L^{Zd} + \frac{3g_L^2 - g_Y^2}{4g_Y^2} \delta g_R^{Zu} + \frac{3g_L^2 + g_Y^2}{2g_Y^2} \delta g_R^{Zd}$$

- It is useful to rearrange these 4 couplings so that we can separate the blind direction from the rest of the parameter space:

$$\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = R \begin{pmatrix} \delta g_L^{Zu} \\ \delta g_R^{Zu} \\ \delta g_L^{Zd} \\ \delta g_R^{Zd} \end{pmatrix} = \begin{pmatrix} 0.93 & -0.29 & -0.23 & -0.01 \\ 0.18 & 0.87 & -0.33 & -0.33 \\ 0.27 & 0.18 & 0.90 & -0.29 \\ 0.17 & 0.37 & 0.17 & 0.90 \end{pmatrix} \begin{pmatrix} \delta g_L^{Zu} \\ \delta g_R^{Zu} \\ \delta g_L^{Zd} \\ \delta g_R^{Zd} \end{pmatrix}$$



$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.9 \pm 1.8 \\ 0.3 \pm 3.3 \\ -2.4 \pm 4.8 \end{pmatrix} \times 10^{-2}$$

This can be achieved using D0 data [Efrati, Falkowski, Soreq, ‘15] but with very modest precision: $|t| < 0.2$

- t unconstrained.** Can we use LHC data to restrict it?


4. The A_{FB} asymmetry at the LHC

- We find that the cleanest observable for the task at hand is the Drell-Yan forward-backward asymmetry (A_{FB})

$$\frac{d\sigma_{pp}(Y, \hat{s}, \cos \theta^*)}{dY d\hat{s} d\cos \theta^*} \propto \sum_{q=u,d,s,c,b} \left[\hat{\sigma}_{q\bar{q}}^{even}(\hat{s}, \cos \theta^*) + D_{q\bar{q}}(Y, \hat{s}) \hat{\sigma}_{q\bar{q}}^{odd}(\hat{s}, \cos \theta^*) \right] F_{q\bar{q}}(Y, \hat{s})$$

Forward events: $\cos \theta^* > 0$

Backward events: $\cos \theta^* < 0$




4. The A_{FB} asymmetry at the LHC

- We find that the cleanest observable for the task at hand is the Drell-Yan forward-backward asymmetry (A_{FB})

$$\frac{d\sigma_{pp}(Y, \hat{s}, \cos \theta^*)}{dY d\hat{s} d\cos \theta^*} \propto \sum_{q=u,d,s,c,b} \left[\hat{\sigma}_{q\bar{q}}^{even}(\hat{s}, \cos \theta^*) + D_{q\bar{q}}(Y, \hat{s}) \hat{\sigma}_{q\bar{q}}^{odd}(\hat{s}, \cos \theta^*) \right] F_{q\bar{q}}(Y, \hat{s})$$

$f(Zqq)$

Forward events: $\cos \theta^* > 0$
 Backward events: $\cos \theta^* < 0$



4. The A_{FB} asymmetry at the LHC

- We find that the cleanest observable for the task at hand is the Drell-Yan forward-backward asymmetry (A_{FB})

$$\frac{d\sigma_{pp}(Y, \hat{s}, \cos \theta^*)}{dY d\hat{s} d\cos \theta^*} \propto \sum_{q=u,d,s,c,b} \left[\hat{\sigma}_{q\bar{q}}^{even}(\hat{s}, \cos \theta^*) + D_{q\bar{q}}(Y, \hat{s}) \hat{\sigma}_{q\bar{q}}^{odd}(\hat{s}, \cos \theta^*) \right] F_{q\bar{q}}(Y, \hat{s})$$

$f(Zqq)$

$$A_{FB}(Y, \hat{s}) = \frac{\sigma_F(Y, \hat{s}) - \sigma_B(Y, \hat{s})}{\sigma_F(Y, \hat{s}) + \sigma_B(Y, \hat{s})} = SM(1 + \#\delta g_i + \dots)$$

4. The A_{FB} asymmetry at the LHC

$ Y $	Experimental value	SM prediction
0.0 - 0.8	0.0195 ± 0.0015	0.0144 ± 0.0007
0.8 - 1.6	0.0448 ± 0.0016	0.0471 ± 0.0017
1.6 - 2.5	0.0923 ± 0.0026	0.0928 ± 0.0021
2.5 - 3.6	0.1445 ± 0.0046	0.1464 ± 0.0021

Exp. value: [ATLAS-CONF-2018-037 (2018)]

NNLO in QCD SM prediction:

[Bozzi et al., 1007.2351;

Catani et al., 0903.2120;

Catani et al., 1507.06937]

$$A_4 = (3/8)A_{FB}$$

4. The A_{FB} asymmetry at the LHC

$ Y $	Experimental value	SM prediction
0.0 - 0.8	0.0195 ± 0.0015	0.0144 ± 0.0007
0.8 - 1.6	0.0448 ± 0.0016	0.0471 ± 0.0017
1.6 - 2.5	0.0923 ± 0.0026	0.0928 ± 0.0021
2.5 - 3.6	0.1445 ± 0.0046	0.1464 ± 0.0021

Exp. value: [ATLAS-CONF-2018-037 (2018)]

NNLO in QCD SM prediction:

[Bozzi et al., 1007.2351;

Catani et al., 0903.2120;

Catani et al., 1507.06937]

$$A_4 = (3/8)A_{FB}$$

○ Restrictions from each bin:

$$0.0 < |Y| < 0.8 : \quad 0.63 \delta g_L^{Zu} + 0.71 \delta g_R^{Zu} - 0.20 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = 0.088(29)$$

$$0.8 < |Y| < 1.6 : \quad 0.60 \delta g_L^{Zu} + 0.74 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = -0.012(12)$$

$$1.6 < |Y| < 2.5 : \quad 0.53 \delta g_L^{Zu} + 0.80 \delta g_R^{Zu} - 0.16 \delta g_L^{Zd} - 0.23 \delta g_R^{Zd} = -0.0014(92)$$

$$2.5 < |Y| < 3.6 : \quad 0.43 \delta g_L^{Zu} + 0.86 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.21 \delta g_R^{Zd} = -0.0030(81)$$

4. The A_{FB} asymmetry at the LHC

$ Y $	Experimental value	SM prediction
0.0 - 0.8	0.0195 ± 0.0015	0.0144 ± 0.0007
0.8 - 1.6	0.0448 ± 0.0016	0.0471 ± 0.0017
1.6 - 2.5	0.0923 ± 0.0026	0.0928 ± 0.0021
2.5 - 3.6	0.1445 ± 0.0046	0.1464 ± 0.0021

Exp. value: [ATLAS-CONF-2018-037 (2018)]

NNLO in QCD SM prediction:

[Bozzi et al., 1007.2351;

Catani et al., 0903.2120;

Catani et al., 1507.06937]

$$A_4 = (3/8)A_{FB}$$

○ Restrictions from each bin:

$$0.0 < |Y| < 0.8 : \quad 0.63 \delta g_L^{Zu} + 0.71 \delta g_R^{Zu} - 0.20 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = 0.088(29)$$

$$0.8 < |Y| < 1.6 : \quad 0.60 \delta g_L^{Zu} + 0.74 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = -0.012(12)$$

$$1.6 < |Y| < 2.5 : \quad 0.53 \delta g_L^{Zu} + 0.80 \delta g_R^{Zu} - 0.16 \delta g_L^{Zd} - 0.23 \delta g_R^{Zd} = -0.0014(92)$$

$$2.5 < |Y| < 3.6 : \quad 0.43 \delta g_L^{Zu} + 0.86 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.21 \delta g_R^{Zd} = -0.0030(81)$$



○ Restrictions on the four uncorrelated and orthonormal linear combinations:

$$\begin{pmatrix} x' = 0.21 \delta g_L^{Zu} + 0.19 \delta g_R^{Zu} + 0.46 \delta g_L^{Zd} + 0.84 \delta g_R^{Zd} \\ y' = 0.03 \delta g_L^{Zu} - 0.07 \delta g_R^{Zu} - 0.87 \delta g_L^{Zd} + 0.49 \delta g_R^{Zd} \\ z' = 0.83 \delta g_L^{Zu} - 0.54 \delta g_R^{Zu} + 0.02 \delta g_L^{Zd} - 0.10 \delta g_R^{Zd} \\ t' = 0.51 \delta g_L^{Zu} + 0.82 \delta g_R^{Zu} - 0.17 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} \end{pmatrix} = \begin{pmatrix} -10 \pm 4 \\ 0.5 \pm 0.4 \\ 0.04 \pm 0.06 \\ -0.001 \pm 0.005 \end{pmatrix}$$

4. The A_{FB} asymmetry at the LHC

$ Y $	Experimental value	SM prediction
0.0 - 0.8	0.0195 ± 0.0015	0.0144 ± 0.0007
0.8 - 1.6	0.0448 ± 0.0016	0.0471 ± 0.0017
1.6 - 2.5	0.0923 ± 0.0026	0.0928 ± 0.0021
2.5 - 3.6	0.1445 ± 0.0046	0.1464 ± 0.0021

Exp. value: [ATLAS-CONF-2018-037 (2018)]

NNLO in QCD SM prediction:

[Bozzi et al., 1007.2351;

Catani et al., 0903.2120;

Catani et al., 1507.06937]

$$A_4 = (3/8)A_{FB}$$

○ Restrictions from each bin:

$$0.0 < |Y| < 0.8 : \quad 0.63 \delta g_L^{Zu} + 0.71 \delta g_R^{Zu} - 0.20 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = 0.088(29)$$

$$0.8 < |Y| < 1.6 : \quad 0.60 \delta g_L^{Zu} + 0.74 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.22 \delta g_R^{Zd} = -0.012(12)$$

$$1.6 < |Y| < 2.5 : \quad 0.53 \delta g_L^{Zu} + 0.80 \delta g_R^{Zu} - 0.16 \delta g_L^{Zd} - 0.23 \delta g_R^{Zd} = -0.0014(92)$$

$$2.5 < |Y| < 3.6 : \quad 0.43 \delta g_L^{Zu} + 0.86 \delta g_R^{Zu} - 0.18 \delta g_L^{Zd} - 0.21 \delta g_R^{Zd} = -0.0030(81)$$



○ Restrictions on the four uncorrelated and orthonormal linear combinations:


$$\begin{pmatrix} x' = 0.21\delta g_L^{Zu} + 0.19\delta g_R^{Zu} + 0.46\delta g_L^{Zd} + 0.84\delta g_R^{Zd} \\ y' = 0.03\delta g_L^{Zu} - 0.07\delta g_R^{Zu} - 0.87\delta g_L^{Zd} + 0.49\delta g_R^{Zd} \\ z' = 0.83\delta g_L^{Zu} - 0.54\delta g_R^{Zu} + 0.02\delta g_L^{Zd} - 0.10\delta g_R^{Zd} \\ t' = 0.51\delta g_L^{Zu} + 0.82\delta g_R^{Zu} - 0.17\delta g_L^{Zd} - 0.22\delta g_R^{Zd} \end{pmatrix} = \begin{pmatrix} -10 \pm 4 \\ 0.5 \pm 0.4 \\ 0.04 \pm 0.06 \\ -0.001 \pm 0.005 \end{pmatrix}$$

We are capable of obtaining per mille level constraints

4. The A_{FB} asymmetry at the LHC

- Impact on the global fit:

$$\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 0.004 \pm 0.017 \\ 0.010 \pm 0.032 \\ 0.021 \pm 0.046 \\ -0.03 \pm 0.19 \end{pmatrix}, \quad \rho = \begin{pmatrix} 1. & -0.09 & -0.08 & -0.04 \\ -0.09 & 1. & -0.09 & -0.93 \\ -0.08 & -0.09 & 1. & -0.19 \\ -0.04 & -0.93 & -0.19 & 1. \end{pmatrix}$$




The combination of LEP+LHC is good enough to lift the blind direction, but we are not as restrictive as in t' , since $t \cdot t' = 0.16$

4. The A_{FB} asymmetry at the LHC

- Impact on the global fit:

$$\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 0.004 \pm 0.017 \\ 0.010 \pm 0.032 \\ 0.021 \pm 0.046 \\ -0.03 \pm 0.19 \end{pmatrix}, \quad \rho = \begin{pmatrix} 1. & -0.09 & -0.08 & -0.04 \\ -0.09 & 1. & -0.09 & -0.93 \\ -0.08 & -0.09 & 1. & -0.19 \\ -0.04 & -0.93 & -0.19 & 1. \end{pmatrix}$$



The combination of LEP+LHC is good enough to lift the blind direction, but we are not as restrictive as in t' , since $t \cdot t' = 0.16$


- LHC constrains a specific direction much strongly than D0. Both hadron measurements are important for the global fit, although for simple scenarios LHC has a larger effect. All in all, **“traditional pole” observables + ATLAS + D0** give:

$$\begin{pmatrix} \delta g_L^{Zu} \\ \delta g_R^{Zu} \\ \delta g_L^{Zd} \\ \delta g_R^{Zd} \end{pmatrix} = \begin{pmatrix} -0.012 \pm 0.024 \\ -0.005 \pm 0.032 \\ -0.020 \pm 0.037 \\ -0.03 \pm 0.13 \end{pmatrix}, \quad \rho = \begin{pmatrix} 1 & 0.51 & 0.68 & 0.69 \\ 0.51 & 1 & 0.56 & 0.94 \\ 0.68 & 0.56 & 1 & 0.54 \\ 0.69 & 0.94 & 0.54 & 1 \end{pmatrix}$$

The other 16 parameters are also being fitted here, to almost no changes in their limits


4. The A_{FB} asymmetry at the LHC

- A_{FB}^{LHC} provides crucial information in simple NP scenarios:




4. The A_{FB} asymmetry at the LHC

- The use of these two inputs leaves much less room for the inclusion of nonlinear contributions:



4. The A_{FB} asymmetry at the LHC

- The use of these two inputs leaves much less room for the inclusion of nonlinear contributions:



5. Conclusions

- LHC A_{FB} provides $\sim 0.5\%$ bounds on Zqq corrections

$$0.51\delta g_L^{Zu} + 0.82\delta g_R^{Zu} - 0.17\delta g_L^{Zd} - 0.22\delta g_R^{Zd} = -0.001 \pm 0.005$$

- **The t variable is lifted with the inclusion of the A_{FB} ATLAS input**
 - We find that the ATLAS A_{FB} information provides a significant improvement on LEP-only bounds on the Zqq vertex corrections even in simple scenarios with few free parameters
-

5. Conclusions

- LHC A_{FB} provides $\sim 0.5\%$ bounds on Zqq corrections


$$0.51\delta g_L^{Zu} + 0.82\delta g_R^{Zu} - 0.17\delta g_L^{Zd} - 0.22\delta g_R^{Zd} = -0.001 \pm 0.005$$

- **The t variable is lifted with the inclusion of the A_{FB} ATLAS input**
- We find that the ATLAS A_{FB} information provides a significant improvement on LEP-only bounds on the Zqq vertex corrections even in simple scenarios with few free parameters

-
- **Outlook 1:** Current and future measurements of Drell-Yan dilepton production by LHC could be analyzed following a similar procedure to ours in order to extend the impact of hadron colliders on the electroweak precision program
 - **Outlook 2:** Information from Drell-Yan cross sections could be added, and off-pole data could be analyzed too (\rightarrow LLQQ operators enter)

EXTRA SLIDES

Backup 1: A_{FB} impact on the global SMEFT fit



Backup 2: Allowed regions for some simple NP settings

