

Semi-leptonic τ decays BSM

[Based on: V. Cirigliano, D. Díaz-Calderón, A. Falkowski, M. González-Alonso, & A. Rodríguez-Sánchez, arxiv:2112.02087]

David Díaz Calderón

IFIC (CSIC-UV)



VNIVERSITAT
ID VALÈNCIA



Overview

- Introduction.
- Theoretical framework.
- Hadronic tau decays
- Combination of bounds.
- Other probes
 - π decays.
 - K + Hyperon decays.
 - Nuclear β decays.
- Global fit.
- Conclusions

Introduction

Great experimental and theoretical precision in hadronic tau decays:

- α_S , V_{US} , f_π , QCD vacuum condensates, ...

Introduction

Great experimental and theoretical precision in hadronic tau decays:


- α_S , V_{US} , f_π , QCD vacuum condensates, ...

Tension with other determinations

Introduction

Great experimental and theoretical precision in hadronic tau decays:


- α_S , V_{US} , f_π , QCD vacuum condensates, ...



Introduction

Great experimental and theoretical precision in hadronic tau decays:


- α_S , V_{US} , f_π , QCD vacuum condensates, ...



Introduction

Great experimental and theoretical precision in hadronic tau decays:

- α_S , V_{US} , f_π , QCD vacuum condensates, ...



Theoretical Framework


$$\begin{aligned}\mathcal{L}_{\text{eff}} = & -\frac{G_\mu V_{uD}}{\sqrt{2}} \left[\left(1 + \epsilon_L^{D\ell}\right) \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 - \gamma_5) D \right. \\ & + \epsilon_R^{D\ell} \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 + \gamma_5) D + \bar{\ell} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \left[\epsilon_S^{D\ell} - \epsilon_P^{D\ell} \gamma_5 \right] D \\ & \left. + \frac{1}{4} \hat{\epsilon}_T^{D\ell} \bar{\ell} \sigma_{\mu\nu} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \sigma^{\mu\nu} (1 - \gamma_5) D \right] + \text{h.c.}\end{aligned}$$

Theoretical Framework

$$\begin{aligned}
 \mathcal{L}_{\text{eff}} = & -\frac{G_\mu V_{uD}}{\sqrt{2}} \left[\left(1 + \epsilon_L^{D\ell}\right) \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 - \gamma_5) D \right. \\
 & + \epsilon_R^{D\ell} \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 + \gamma_5) D + \bar{\ell} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \left[\epsilon_S^{D\ell} - \epsilon_P^{D\ell} \gamma_5 \right] D \\
 & \left. + \frac{1}{4} \epsilon_T^{D\ell} \bar{\ell} \sigma_{\mu\nu} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \sigma^{\mu\nu} (1 - \gamma_5) D \right] + \text{h.c.}
 \end{aligned}$$



Theoretical Framework

$$\begin{aligned}
 \mathcal{L}_{\text{eff}} = & -\frac{G_\mu V_{uD}}{\sqrt{2}} \left[\left(1 + \epsilon_L^{D\ell}\right) \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 - \gamma_5) D \right. \\
 & + \epsilon_R^{D\ell} \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 + \gamma_5) D + \bar{\ell} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \left[\epsilon_S^{D\ell} - \epsilon_P^{D\ell} \gamma_5 \right] D \\
 & \left. + \frac{1}{4} \epsilon_T^{D\ell} \bar{\ell} \sigma_{\mu\nu} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \sigma^{\mu\nu} (1 - \gamma_5) D \right] + \text{h.c.}
 \end{aligned}$$



Theoretical Framework

$$\mathcal{L}_{\text{eff}} = -\frac{G_\mu V_{uD}}{\sqrt{2}} \left[\left(1 + \epsilon_L^{D\ell}\right) \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 - \gamma_5) D \right. \\ \left. + \epsilon_R^{D\ell} \bar{\ell} \gamma_\mu (1 - \gamma_5) \nu_\ell \cdot \bar{u} \gamma^\mu (1 + \gamma_5) D + \bar{\ell} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \left[\epsilon_S^{D\ell} - \epsilon_P^{D\ell} \gamma_5 \right] D \right. \\ \left. + \frac{1}{4} \epsilon_T^{D\ell} \bar{\ell} \sigma_{\mu\nu} (1 - \gamma_5) \nu_\ell \cdot \bar{u} \sigma^{\mu\nu} (1 - \gamma_5) D \right] + \text{h.c.}$$



Hadronic τ Decays

Hadronic τ Decays

$$\tau \rightarrow \pi\nu, \tau \rightarrow K\nu$$

$$\tau \rightarrow \pi\pi\nu$$

$$\tau \rightarrow \eta\pi\nu$$

Non-strange inclusive

Strange inclusive

Hadronic τ Decays

$\tau \rightarrow \pi\nu, \tau \rightarrow K\nu \longrightarrow \epsilon_L^{D\tau} - \epsilon_L^{De}, \epsilon_R^D$ and $\epsilon_P^{D\tau}$.

$\tau \rightarrow \pi\pi\nu \longrightarrow \epsilon_L^{d\tau} - \epsilon_L^{de}$ and $\epsilon_T^{d\tau}$. $\epsilon_S^{d\tau}$ is suppressed.

$\tau \rightarrow \eta\pi\nu \longrightarrow \epsilon_S^{d\tau}$ enhanced \rightarrow only constrains $\epsilon_S^{d\tau}$.

Non-strange inclusive \longrightarrow Isospin Symmetry $\rightarrow \epsilon_L^{d\tau} - \epsilon_L^{de}, \epsilon_R^{d\tau}$ and $\epsilon_T^{d\tau}$.

Strange inclusive \longrightarrow SU(3) $\rightarrow \epsilon_L^{s\tau} - \epsilon_L^{se}, \epsilon_R^{s\tau}, \epsilon_T^{s\tau}, \epsilon_S^{s\tau}$ and $\epsilon_P^{s\tau}$.

Hadronic τ Decays: constrains

$$\tau \rightarrow \pi \nu \xrightarrow{\Gamma(\tau \rightarrow \pi \nu)} \epsilon_L^{d\tau} - \epsilon_L^{de} - \epsilon_R^{d\tau} - \epsilon_R^{de} - \frac{B_0^d}{m_\tau} \epsilon_P^{d\tau} = -(0.9 \pm 7.3) \times 10^{-3}$$

$$\tau \rightarrow K \nu \xrightarrow{\Gamma(\tau \rightarrow K \nu)} \epsilon_L^{s\tau} - \epsilon_L^{se} - \epsilon_R^{s\tau} - \epsilon_R^{se} - \frac{B_0^s}{m_\tau} \epsilon_P^{s\tau} = -(2 \pm 10) \times 10^{-3}$$

$$\tau \rightarrow \pi \pi \nu \xrightarrow{a_\mu^{\text{had, LO}}} \epsilon_L^{d\tau} - \epsilon_L^{de} + \epsilon_R^{d\tau} - \epsilon_R^{de} + 0.43(8) \hat{\epsilon}_T^{d\tau} = (10.0 \pm 4.9) \times 10^{-3}$$

$$\tau \rightarrow \eta \pi \nu \xrightarrow{\text{BR}(\tau \rightarrow \eta \pi \nu)} \epsilon_S^{d\tau} \in (-0.021, 0.0010), \quad |\text{Im}(\epsilon_S^{d\tau})| < 0.014$$

Hadronic τ Decays: constrains

$$\left. \begin{aligned}
 \epsilon_{L+R}^{d\tau} - \epsilon_{L+R}^{de} - 0.76\epsilon_R^{d\tau} + 0.49(16)\hat{\epsilon}_T^{d\tau} &= (4 \pm 10) \times 10^{-3} \\
 \epsilon_{L+R}^{d\tau} - \epsilon_{L+R}^{de} - 0.88\epsilon_R^{d\tau} + 0.27(9)\hat{\epsilon}_T^{d\tau} &= (9.1 \pm 8.8) \times 10^{-3} \\
 \epsilon_{L+R}^{d\tau} - \epsilon_{L+R}^{de} + 3.05\epsilon_R^{d\tau} + 1.9(1.2)\hat{\epsilon}_T^{d\tau} &= (5 \pm 51) \times 10^{-3} \\
 \epsilon_{L+R}^{d\tau} - \epsilon_{L+R}^{de} + 1.93\epsilon_R^{d\tau} + 1.6(1.5)\hat{\epsilon}_T^{d\tau} &= (7.0 \pm 9.5) \times 10^{-3}
 \end{aligned} \right\} \begin{array}{l} \rho_{V+A} \\ \rho_{V-A} \end{array} \left. \vphantom{\begin{array}{l} \rho_{V+A} \\ \rho_{V-A} \end{array}} \right\} \text{Non-strange Inclusive}$$

$$\begin{aligned}
 &1.00 (\epsilon_{L+R}^{s\tau} - \epsilon_{L+R}^{se}) - 1.03 \epsilon_R^{s\tau} - 0.38 \epsilon_P^{s\tau} + 0.40(13) \hat{\epsilon}_T^{s\tau} + 0.08(1) \epsilon_S^{s\tau} \\
 &- 1.07 (\epsilon_{L+R}^{d\tau} - \epsilon_{L+R}^{de}) + 1.04 \epsilon_R^{d\tau} + 0.30 \epsilon_P^{d\tau} - 0.43(14) \hat{\epsilon}_T^{d\tau} \\
 &= -(0.0171 \pm 0.0085)
 \end{aligned} \left. \vphantom{\begin{array}{l} \epsilon_{L+R}^{s\tau} \\ \epsilon_{L+R}^{de} \\ \epsilon_R^{d\tau} \\ \epsilon_P^{d\tau} \\ \hat{\epsilon}_T^{d\tau} \end{array}} \right\} \text{Strange Inclusive}$$

$$|\widehat{V}_{us}|^{\text{inc}} = \left(\frac{\hat{R}_\tau^s}{\hat{R}_\tau^d / |\hat{V}_{ud}|^2 - \delta R_{\text{th}}^{\text{SM}}} \right)^{1/2}$$

Hadronic τ Decays: constrains

	$\epsilon_L^{d\tau} \times 10^3$	$\epsilon_L^{de} \times 10^3$	$\epsilon_R^d \times 10^3$	$\epsilon_P^{d\tau} \times 10^3$	$\epsilon_T^{d\tau} \times 10^3$	$\epsilon_S^{d\tau} \times 10^3$
$\tau \rightarrow \pi\nu$	-0.9(7.3)	0.9(7.3)	0.9(7.3)	0.6(5.0)	x	x
$\tau \rightarrow \pi\pi\nu$	10(4.9)	-10(4.9)	x	x	23(12)	x
$\tau \rightarrow \pi\eta\nu$	x	x	x	x	x	(-21, 10)
$V + A$	6.9(7.0)	-6.9(7.0)	-8.6(8.4)	x	15(19)	x
$V - A$	7.0(9.5)	-7.0(9.5)	3.6(4.9)	x	15(17)	x
	$\epsilon_L^{s\tau} \times 10^3$	$\epsilon_L^{se} \times 10^3$	$\epsilon_R^s \times 10^3$	$\epsilon_P^{s\tau} \times 10^3$	$\epsilon_T^{s\tau} \times 10^3$	$\epsilon_S^{s\tau} \times 10^3$
$\tau \rightarrow K\nu$	-2(10)	2(10)	2(10)	1.2(6.1)	x	x
S. Inclusive	-17(16)	17(16)	23(22)	340(327)	-34(35)	-170(161)

Hadronic τ Decays: fit

$$\begin{pmatrix} \epsilon_L^{d\tau/e} + \epsilon_R^{d\tau} - \epsilon_R^{de} \\ \epsilon_R^{d\tau} \\ \epsilon_P^{d\tau} \\ \hat{\epsilon}_T^{d\tau} \\ \epsilon_L^{s\tau/e} - \epsilon_R^{s\tau} - \epsilon_R^{se} - \frac{m_{K^\pm}^2}{m_\tau(m_u+m_s)}\epsilon_P^{s\tau} \\ \epsilon_L^{s\tau/e} - 0.03\epsilon_R^{s\tau} - \epsilon_R^{se} + 0.08(1)\epsilon_S^{s\tau} - 0.38\epsilon_P^{s\tau} + 0.40(13)\hat{\epsilon}_T^{s\tau} \end{pmatrix} = \begin{pmatrix} 2.4 \pm 2.6 \\ 0.7 \pm 1.4 \\ 0.4 \pm 1.0 \\ -3.3 \pm 6.0 \\ -0.2 \pm 1.0 \\ -1.3 \pm 1.2 \end{pmatrix} \times 10^{-2},$$

$$\left(\epsilon_L^{D\tau/e} \equiv \epsilon_L^{D\tau} - \epsilon_L^{De} \right)$$

$$\rho = \begin{pmatrix} 1 & 0.87 & -0.18 & -0.98 & -0.03 & -0.45 \\ & 1 & -0.59 & -0.86 & 0.06 & -0.59 \\ & & 1 & 0.18 & -0.36 & 0.38 \\ & & & 1 & 0.04 & 0.49 \\ & & & & 1 & 0.16 \\ & & & & & 1 \end{pmatrix}.$$

→ Percent level marginalized constrains.

→ All Lorentz structures resolved in the $d\tau$ sector.

→ Only two combinations of $\epsilon_X^{s\tau}$ are constrained.




We cannot resolve $\epsilon_X^{s\tau}$

Other probes

nuclear + π decays


K decay + Hyperon β decay

Other probes



K decay + Hyperon β decay

Other probes



Other probes

nuclear + π decays

$$\begin{pmatrix} \hat{V}_{ud} \\ \epsilon_R^d \\ \epsilon_S^{de} \\ \hat{\epsilon}_T^{de} \\ \epsilon_P^{de} \\ \epsilon_L^{d\mu} - \epsilon_L^{de} - \epsilon_P^{d\mu} \frac{m_\pi^2}{m_\mu(m_u+m_d)} \end{pmatrix} = \begin{pmatrix} 0.97386(40) \\ -0.012(12) \\ 0.00032(99) \\ -0.0004(11) \\ 3.9(4.3) \times 10^{-6} \\ -0.021(24) \end{pmatrix}$$

K decay + Hyperon β decay

$$\begin{pmatrix} \hat{V}_{us} \\ \epsilon_L^{s\mu} - \epsilon_L^{se} \\ \epsilon_R^s \\ \epsilon_S^{s\mu} \\ \epsilon_P^{se} \\ \epsilon_P^{s\mu} \\ \hat{\epsilon}_T^{s\mu} \end{pmatrix} = \begin{pmatrix} 0.22306(56) \\ 0.0008(22) \\ 0.001(50) \\ -0.00026(44) \\ -0.3(2.0) \times 10^{-5} \\ -0.0006(41) \\ 0.002(22) \end{pmatrix}$$

Global fit


$$\begin{pmatrix}
 \hat{V}_{us} \equiv V_{us}(1 + \epsilon_L^{se} + \epsilon_R^s) \\
 \epsilon_L^{dse} \equiv \epsilon_L^{de} + \frac{\hat{V}_{us}^2}{1 - \hat{V}_{us}^2} \epsilon_L^{se} \\
 \epsilon_R^d \\
 \epsilon_S^{de} \\
 \epsilon_P^{de} \\
 \hat{\epsilon}_T^{de} \\
 \epsilon_L^{s\mu} - \epsilon_L^{se} \\
 \epsilon_R^s \\
 \epsilon_P^{se} \\
 \epsilon_L^{d\mu} - \epsilon_L^{de} - \epsilon_P^{d\mu} \frac{m_{\pi^\pm}^2}{m_\mu(m_u + m_d)} \\
 \epsilon_S^{s\mu} \\
 \epsilon_P^{s\mu} \\
 \hat{\epsilon}_T^{s\mu} \\
 \epsilon_L^{d\tau} - \epsilon_L^{de} \\
 \epsilon_P^{d\tau} \\
 \hat{\epsilon}_T^{d\tau} \\
 \epsilon_L^{s\tau} - \epsilon_L^{se} - \epsilon_P^{s\tau} \frac{m_{K^\pm}^2}{m_\tau(m_u + m_s)} \\
 \epsilon_L^{s\tau} - \epsilon_L^{se} + 0.08(1)\epsilon_S^{s\tau} - 0.38\epsilon_P^{s\tau} + 0.40(13)\hat{\epsilon}_T^{s\tau}
 \end{pmatrix}
 =
 \begin{pmatrix}
 0.22306(56) \\
 2.2(8.6) \\
 -3.3(8.2) \\
 3.0(9.9) \\
 1.3(3.4) \\
 -0.4(1.1) \\
 0.8(2.2) \\
 0.2(5.0) \\
 -0.3(2.0) \\
 -0.5(1.8) \\
 -2.6(4.4) \\
 -0.6(4.1) \\
 0.2(2.2) \\
 0.1(1.9) \\
 9.2(8.6) \\
 1.9(4.5) \\
 0.0(1.0) \\
 -0.7(5.2)
 \end{pmatrix}
 \times 10^\wedge
 \begin{pmatrix}
 0 \\
 -3 \\
 -3 \\
 -4 \\
 -6 \\
 -3 \\
 -3 \\
 -2 \\
 -5 \\
 -2 \\
 -4 \\
 -3 \\
 -2 \\
 -2 \\
 -3 \\
 -2 \\
 -1 \\
 -2
 \end{pmatrix}$$

$$\chi_{SM}^2 - \chi_{min}^2 = 37.4 \Rightarrow 3\sigma$$

Global fit

$$\chi_{SM}^2 - \chi_{min}^2 = 37.4 \Rightarrow 3\sigma$$

Why?




Cabibbo anomaly \rightarrow Inconsistency in V_{us} determinations

Global fit

$$\chi_{SM}^2 - \chi_{min}^2 = 37.4 \Rightarrow 3\sigma$$

Why?



Cabibbo anomaly \rightarrow Inconsistency in V_{us} determinations

Simple BSM models can fix the anomaly

Global fit


One-at-a-time fit

	$\epsilon_X^{de} \times 10^3$	$\epsilon_X^{se} \times 10^3$	$\epsilon_X^{d\mu} \times 10^3$	$\epsilon_X^{s\mu} \times 10^3$	$\epsilon_X^{d\tau} \times 10^3$	$\epsilon_X^{s\tau} \times 10^3$
L	-0.79(25)	-0.6(1.2)	0.40(87)	0.5(1.2)	5.0(2.5)	-18.2(6.2)
R	-0.62(25)	-5.2(1.7)	-0.62(25)	-5.2(1.7)	-0.62(25)	-5.2(1.7)
S	1.40(65)	-1.6(3.2)	x	-0.51(43)	-6(16)	-270(100)
P	0.00018(17)	-0.00044(36)	-0.015(32)	-0.032(64)	1.7(2.5)	10.4(5.5)
\hat{T}	0.29(82)	0.035(70)	x	2(18)	28(10)	-55(27)

In red: 3σ or more preference for BSM

→ $\epsilon_R^s, \epsilon_L^{de}$ ease the tension between nuclear and kaon decays.

→ $\epsilon_L^{s\tau}$ eases the tension between $\tau \rightarrow s$ inclusive and kaon decays.



$$\epsilon_R^d, \epsilon_R^s, \epsilon_L^{s\tau}$$

$$\chi_{\text{SM}}^2 - \chi_{\text{min}}^2 = 26.1 \Rightarrow 4.4\sigma$$

Conclusions

- Most comprehensive discussion of multi-body and inclusive hadronic τ decays.

Conclusions

- Most comprehensive discussion of multi-body and inclusive hadronic τ decays.
- We provided a model-independent global analysis of semileptonic charged-current decays of light quarks involving all three lepton families.

Conclusions

- Most comprehensive discussion of multi-body and inclusive hadronic τ decays.
- We provided a model-independent global analysis of semileptonic charged-current decays of light quarks involving all three lepton families.
- Hadronic τ decays do not show a particular preference for new physics.

Conclusions

- Most comprehensive discussion of multi-body and inclusive hadronic τ decays.
- We provided a model-independent global analysis of semileptonic charged-current decays of light quarks involving all three lepton families.
- Hadronic τ decays do not show a particular preference for new physics.
- This changes when nuclear β , baryon, pion and kaon decays are included \rightarrow Cabibbo anomalies.

Conclusions

- Most comprehensive discussion of multi-body and inclusive hadronic τ decays.
- We provided a model-independent global analysis of semileptonic charged-current decays of light quarks involving all three lepton families.
- Hadronic τ decays do not show a particular preference for new physics.
- This changes when nuclear β , baryon, pion and kaon decays are included \rightarrow Cabibbo anomalies.
- These anomalies can be eased in simple BSM scenarios.