

Octupole correlations in the neutron-deficient ^{110}Xe nucleus

Thursday, 30 May 2024 12:50 (20)

Octupole correlations near $N = Z = 56$ are unique in sense that they occur between particles in the same orbitals for both neutrons and protons. In this region just above ^{100}Sn , it is expected that enhanced octupole correlations will take place at low and medium spins in the light Te ($Z = 52$), I ($Z = 53$) and Xe ($Z = 54$) nuclei [1]. In this region of the nuclear chart, the Fermi surface for both neutrons and protons lies close to orbitals from the $d_{5/2}$ and $h_{11/2}$ subshells; octupole correlations emerge from the interactions of particles in these orbitals with valence neutrons and protons outside the ^{100}Sn core [2, 3]. As a result of the octupole correlations, an enhancement of octupole collectivity is expected to appear. Close to $N = Z = 56$, a level structure characteristic of octupole correlations, consisting of negative-parity states and enhanced E1 transitions, has been observed in a number of cases including ^{112}Xe [4], ^{114}Xe [5, 6, 7] and ^{118}Ba [8].

With the aim to observe for the first time the octupole band in the neutron-deficient ($N = Z + 2$) ^{110}Xe nucleus, an in-beam experiment was performed at the Accelerator Laboratory of the University of Jyväskylä, Finland. The ^{110}Xe nuclei were produced via the $^{54}\text{Fe}(^{58}\text{Ni}, 2n)$ fusion-evaporation reaction. The emitted γ rays were detected using the JUROGAM3 γ -ray spectrometer [9], while the fusion-evaporation residues were separated with the MARA separator [10]. In this experiment, we were able to prove the existence of the octupole band via the identification of the low-lying 3^- and 5^- states, and their inter-band E1 transitions between the ground-state band and the octupole band [11]. Hence, this new experimental findings will be presented combined with a detailed study of the systematics of the energy levels and the $B(E2)/B(E1)$ ratios in $^{110-114}\text{Xe}$, and a comparison with state-of-the-art theoretical calculations.

- [1] G. de Angelis et al., Phys. Lett. B 437 (1998) 236.
- [2] P.A. Butler and W. Nazarewicz, Rev. Mod. Phys. 68 (1996) 349.
- [3] L.M. Robledo and G. F. Bertsch, Phys. Rev. C 84 (2011), 054302.
- [4] J.F. Smith et al. Phys. Lett. 523 B, 13 (2001).
- [5] S.L. Rugari et al. Phys. Rev. C 48, 2078 (1993).
- [6] E.S. Paul et al., Nucl. Phys. A673, 31 (2000).
- [7] G. de Angelis et al., Phys. Lett. B 535 (2002) 93.
- [8] J.F. Smith et al., Phys.Rev. C5 7, R1037-R1041 (1998).
- [9] J. Pakarinen et al., Eur.Phys. J. A 56 (2020) 150.
- [10] J. Sarven et al., Nucl. Instr. and Meth. B 266 (2008) 4196-4200.
- [11] A. Illana et al., Phys. Lett. B 848 (2024) 138371.

Primary author(s) : ILLANA SISON, Andres (UCM); PEREZ VIDAL, Rosa Maria (IFIC)

Presenter(s) : ILLANA SISON, Andres (UCM)

Session Classification : Session 12