

First laser-spectroscopy measurements across $N = 32$ in the calcium isotopic chain

Tuesday, 28 May 2024 18:10 (20)

On behalf of the COLLAPS collaboration

Over a decade ago, the first experimental evidence for the $N=32$ sub shell closure in the calcium isotopic chain emerged [1,2]. Subsequent experimental and theoretical investigations have confirmed this finding. However, in laser spectroscopy measurements extending up to ^{52}Ca ($N=32$), no indications of this shell gap were apparent [3]. Crossing the shell gap with laser spectroscopy setups has proved difficult due to the simultaneous requirement of a sensitivity of approximately 10 ions/s and a measurement uncertainty on the order of MHz.

This contribution presents the first laser spectroscopy measurements of ^{53}Ca , facilitated by an extension of the collinear laser spectroscopy technique employed at the COLLAPS setup at ISOLDE/CERN. This technique, termed as *radioactive detection after optical pumping and state selective charge exchange* (ROC), combines the high sensitivity of a particle detection scheme with the high resolution of low-power, continuous wave lasers utilized in a collinear geometry. The methodology of this technique will be explained, followed by the presentation and discussion of preliminary values for the charge radius and magnetic dipole moment of ^{53}Ca in the context of the robustness of the $N=32$ sub shell closure, as well as the prospects to measure ^{54}Ca .

- [1] Wienholtz, F. et al. Nature vol. 498, 346-349 (2013)
- [2] Steppenbeck, D. et al. Nature vol. 502, 207-210 (2013)
- [3] R.F. Garcia Ruiz et al, Nature Physics vol. 12, 594-598 (2016)

Primary author(s) : LELLINGER, Tim (CERN)

Presenter(s) : LELLINGER, Tim (CERN)

Session Classification : Session 8