

LHC differential top-quark pair production cross sections in the ABMP PDF fit

Sergey Alekhin¹, Maria Vittoria Garzelli¹, Sven-Olaf Moch¹, Sasha Zenaiev¹
with input from Javier Mazzitelli²


¹ Hamburg University

² PSI

arxiv:2311.05509 “Top-quark pole mass extraction at NNLO accuracy, from total, single- and double-differential cross sections for $t\bar{t} + X$ production at the LHC” (Garzelli, Mazzitelli, Moch, SZ)
arxiv:24XX.XXXXX in preparation (Alekhin, Garzelli, Moch, SZ)


Workshop on top quark mass measurements
23 May 2024

Why study $t\bar{t}$ production?



- m_t provides a hard scale
 $\Rightarrow$ ultimate probe of pQCD
 (NLO, aNNLO, NNLO, ...)
- Produced mainly via gg
 $\Rightarrow$ constrain gluon PDF at high x
- Production sensitive to α_s and m_t^{pole}
- May provide insight into possible new physics

Example:







- Simultaneous extraction of PDFs, α_s , m_t^{pole} using normalised triple-differential cross sections at NLO
- Extended to $\overline{\text{MS}}$, MSR schemes in JHEP 04 (2021) 043 [Garzelli, Kemmler, Moch, Zenaiev]

Scope of this work

- **NNLO calculations** for total and fully differential $t\bar{t}$ (q_T subtraction) are now publicly available with **MATRIX framework** [Catani et al. Phys.Rev.D 99 (2019) 5, 051501; JHEP 07 (2019) 100]
 - ▶ fully differential NNLO calculations were also published in JHEP 04 (2017) 071 [Czakon, Heymes, Mitov], but no public code available
 - ▶ further details about our MATRIX setup in BACKUP
- **Multi-differential $t\bar{t}$ measurements** enable extraction of **PDFs, α_S , m_t^{pole}**
 - ▶ for 3D x-sections $M(t\bar{t})$, $y(t\bar{t})$, N_{jet} [CMS TOP-18-004] NNLO calculations are not available for $t\bar{t}$ + jets
 - ▶ therefore for the time being we focus on double-differential **$M(t\bar{t})$, $y(t\bar{t})$ x-sections**
 - ★ **$M(t\bar{t})$** provides sensitivity to **m_t**
 - ★ **$y(t\bar{t})$** provides sensitivity to **PDFs** via relation to partonic momentum fraction x :
at LO $x_{1,2} = (M(t\bar{t})/\sqrt{s}) \exp[\pm y(t\bar{t})]$
- We have interfaced MATRIX $t\bar{t}$ to PineAPPL interpolation lib (no NNLO/NLO K-factors etc.)
- Recently we have extracted m_t^{pole} from the total and differential ATLAS and CMS measurements of $t\bar{t}$ production using different PDF sets [Garzelli, Mazzitelli, Moch, Zenaiev arXiv:2311.05509]
 - ▶ overall uncertainty on m_t^{pole} is 0.3 GeV (experimental, PDF, scale variations)
 - ▶ possible extra uncertainty (not accounted for):
 - ★ renormalon ambiguity
 - ★ m_t in MC used to unfold the data to parton level
- **Now we do a global PDF fit with total and differential $t\bar{t}$ within the ABMP16 framework**
 - ▶ using collider and fixed-target DIS, DY, **updated** single top and $t\bar{t}$ data (see BACKUP)

Variation of r cut, validation vs JHEP 04 (2017) 071 by Czakon et al. [CHM]

Good agreement $< 1\%$



Data used in this analysis

Selection of data:


- all measurements of total $\sigma(t\bar{t})$:
 - 10 data points, including recently combined CMS+ATLAS cross section at 7 and 8 TeV
- differential measurements $\frac{1}{\sigma(t\bar{t})} \frac{d\sigma(t\bar{t})}{dO}$ which satisfy following criteria:
 - as function of $M(t\bar{t})$ (if available, 2D $M(t\bar{t})$ and $y(t\bar{t})$)
 - unfolded to parton level (no cuts on p_T , y of leptons or jets): no LHCb data
 - normalized cross sections (to avoid unknown correlation with total $\sigma(t\bar{t})$ and to reduce unknown correlations between different data sets)
 - bin-by-bin correlations are available
 - for the moment only Run-2 2D data included in the PDF fit (besides the total $t\bar{t}$ x-section data)

$\sigma(t\bar{t})$


Experiment	decay channel	dataset	luminosity	$\sqrt{s}$
ATLAS & CMS	combined	2011	5 fb ⁻¹	7 TeV
ATLAS & CMS	combined	2012	20 fb ⁻¹	8 TeV
ATLAS	dileptonic, semileptonic	2011	257 pb ⁻¹	5.02 TeV
CMS	dileptonic	2011	302 pb ⁻¹	5.02 TeV
ATLAS	dileptonic	2015-2018	140 fb ⁻¹	13 TeV
ATLAS	semileptonic	2015-2018	139 fb ⁻¹	13 TeV
CMS	dileptonic	2016	35.9 fb ⁻¹	13 TeV
CMS	semileptonic	2016-2018	137 fb ⁻¹	13 TeV
ATLAS	dileptonic	2022	11.3 fb ⁻¹	13.6 TeV
CMS	dileptonic, semileptonic	2022	1.21 fb ⁻¹	13.6 TeV

$\frac{1}{\sigma(t\bar{t})} \frac{d\sigma(t\bar{t})}{dO}$

Experiment	decay channel	dataset	luminosity	$\sqrt{s}$	observable(s)	n
CMS	semileptonic	2016–2018	137 fb ⁻¹	13 TeV	$M(t\bar{t}), y(t\bar{t}) $	34
CMS	dileptonic	2016	35.9 fb ⁻¹	13 TeV	$M(t\bar{t}), y(t\bar{t}) $	15
ATLAS	semileptonic	2015–2016	36 fb ⁻¹	13 TeV	$M(t\bar{t}), y(t\bar{t}) $	19
ATLAS	all-hadronic	2015–2016	36.1 fb ⁻¹	13 TeV	$M(t\bar{t}), y(t\bar{t}) $	10
CMS	dileptonic	2012	19.7 fb ⁻¹	8 TeV	$M(t\bar{t}), y(t\bar{t}) $	15
ATLAS	semileptonic	2012	20.3 fb ⁻¹	8 TeV	$M(t\bar{t})$	6
ATLAS	dileptonic	2012	20.2 fb ⁻¹	8 TeV	$M(t\bar{t})$	5
ATLAS	dileptonic	2011	4.6 fb ⁻¹	7 TeV	$M(t\bar{t})$	4
ATLAS	semileptonic	2011	4.6 fb ⁻¹	7 TeV	$M(t\bar{t})$	4




CMS 2108.02803 vs NNLO predictions using different PDFs




- Fixed $m_t^{\text{pole}} = 172.5$ GeV (not always consistent with PDF sets), $\mu_r = \mu_f = H_T/4$
- Reported χ^2 values with (and without) PDF uncertainties
- All PDF sets describe data reasonably well, with best description by ABMP16
 - ▶ CT18, MSHT20 and NNPDF40 show clear trend w.r.t data at high $y(t\bar{t})$ (large x)

CMS 2108.02803 vs NNLO predictions using different PDFs



- **NOTE:** m_t^{pole} values on this plot are not the same as the ones obtained in ABMP16 fit
- Low $M(t\bar{t})$: strong dependence on m_t^{pole} via threshold effects
- High $M(t\bar{t})$: opposite dependence due to cross section normalization

CMS 2108.02803 vs NNLO predictions using different PDFs





● ABMP16, fixed $m_t^{\text{pole}} = 172.5 \text{ GeV}$

● Scale variations $< 1\%$ at low $M(t\bar{t})$ (largest cancellation), reach $\approx 4\%$ at high $M(t\bar{t})$

→ these data are useful to provide constraints on m_t and PDFs

Extraction of m_t^{pole} [Garzelli, Mazzitelli, Moch, SZ arXiv:2311.05509]





Data set	n	ABMP16	CT18	MSHT20	NNPDF4.0
CMS 13 TeV semileptonic [48]	34	19(20)	29(176)	38(132)	55(90)
CMS 13 TeV dileptonic [52]	15	15(15)	23(38)	27(34)	23(23)
ATLAS13 TeV semileptonic [53]	19	11(15)	12(17)	11(13)	12(12)
ATLAS 13 TeV all-hadronic [54]	10	11(11)	16(19)	16(17)	14(14)
CMS 8 TeV dileptonic [55]	15	11(15)	11(12)	11(12)	12(12)
ATLAS 8 TeV semileptonic [57]	6	10(12)	4(4)	4(4)	5(5)
ATLAS 7 TeV dileptonic [58]	4	2(3)	1.9(1.9)	1.6(1.6)	1.1(1.1)
ATLAS 8 TeV dileptonic [58]	5	0.2(0.2)	0.4(0.5)	0.4(0.4)	0.2(0.2)
ATLAS 7 TeV semileptonic [56]	4	0.9(1.0)	5(6)	6(6)	3(3)
$\sigma(t\bar{t})$ [43-51]	10	11(26)	16(61)	16(43)	11(12)
Total	122	101(117)	115(337)	113(262)	129(172)

*conservative
uncertainties?*


Discussion of uncertainty treatment

- Experimental uncertainties:
 - ▶ only two data sets (CMS dilepton Run 1, CMS dilepton Run 2) report source by source systematic uncertainties
 - ▶ all other data sets report covariance matrices: it is even not possible to separate systematic correlation from statistical (unfolding)
 - ▶ for some data sets (ATLAS l+jet Run 2, ATLAS dilepton Run 1 (8 TeV)), covariance matrices are not singular as they should be for normalised x-sections (one degree of freedom is lost): numerical inaccuracy?
 - ▶ **we tried to minimize the impact of the lack of experimental systematic correlations by using the total x-section and normalised differential x-sections: many systematic uncertainties cancel for normalised x-sections, but:**
 - please always report covariance matrix from unfolding and source by source systematic uncertainties
 - please specify the sign of systematic variation (e.g. scale varied up or down)
 - if some systematics is calculated as envelope of several variations, please report every variation
 - for covariance matrix, please do some basic consistency checks (positive definiteness, singularity if normalised x-sections etc.)
- Theoretical uncertainties: we assign 1% uncorrelated uncertainty (< data unc.) to cover:
 - ▶ numerical uncertainties of NNLO MATRIX calculations
 - ideally, numerical uncertainty should be stored bin-by-bin in PineAPPL grids, **but storing such uncertainties is not yet possible for any interpolation grids**
 - ▶ small uncertainties due to PineAPPL interpolation tables: negligible (easy to control)
 - ▶ possible systematic differences due to the subtraction scheme (MATRIX vs calculations by Czakon et al.): small, but might become non-negligible in the future

Extraction of m_t^{pole} [Garzelli, Mazzitelli, Moch, SZ arXiv:2311.05509]




CMS ($\sqrt{s}=13$ TeV, 137 fb^{-1} , $pp \rightarrow t\bar{t}X \rightarrow l\text{jet}X$) 2108.02803




- Good description of the data
- Significantly constrained PDF uncertainty band

Other $t\bar{t}$ differential data in ABMP16tt PDF fit


ATLAS ($\sqrt{s}=13$ TeV, 36 fb^{-1} , $pp \rightarrow t\bar{t}X \rightarrow \text{ljetX}$) 1908.07305



ATLAS ($\sqrt{s}=13$ TeV, 36 fb^{-1} , $pp \rightarrow t\bar{t}X \rightarrow \text{hadronsX}$) 2006.09274



CMS ($\sqrt{s}=13$ TeV, 36 fb^{-1} , $pp \rightarrow t\bar{t}X \rightarrow l^+\bar{l}X$) 1904.05237



consistent with
CMS 2402.08486
($\mathcal{L} = 138 \text{ pb}^{-1}$)
see BACKUP

- ATLAS ljet data tend to be **above** theory predictions at $M(t\bar{t}) \gtrsim 1000$ GeV (but still ok within large data uncertainties)
- ATLAS hadronic data tend to be **below** theory predictions at $M(t\bar{t}) \gtrsim 700$ GeV (but still ok within large data uncertainties)
- In summary, all data are in good agreement with NNLO theoretical predictions and put significant constraints on the PDFs

Experiment	Data set	$\sqrt{s}$ (TeV)	Reference	NDP	χ^2		
					I	II	III
ATLAS	$ATLAS\ 13_{ljet}$	13	[25]	19	34.0(42.8)	28.2	–
	$ATLAS\ 13_{had}$	13	[26]	10	11.9(11.7)	11.6	–
CMS	$CMS\ 13_{ll}$	13	[24]	15	20.7(15.9)	–	19.6
	$CMS\ 13_{ljet}$	13	[22]	34	44.3(52.0)	–	42.4


I: both ATLAS and CMS

II: only ATLAS

III: only CMS

→ Overall good description of data by NNLO theoretical predictions, but some tension between ATLAS and CMS differential $t\bar{t}$ data is noticeable

$m_t(m_t)$ and α_s in ABMP16tt PDF fit




Data set	$m_t(m_t)$ (GeV)	χ^2/NDP
<i>ATLAS 13_{ljet}</i>	158.9 ± 1.3	25.2/19
<i>ATLAS 13_{had}</i>	160.5 ± 2.0	11.3/10
<i>CMS 13_{ll}</i>	161.1 ± 1.4	13.9/15
<i>CMS 13_{ljet}</i>	158.7 ± 0.9	37.4/34

- Overall, good agreement between $m_t(m_t)$ extracted from different data sets
- Good agreement with ABMP16 fit and $\sim 50\%$ **reduced uncertainty** on $m_t(m_t)$

$\alpha_s(M_Z, N_f = 5)$		$m_t(m_t)$ (GeV)
Fitted	0.1150(9)	160.6(6)
Fixed	0.114	160.2(4)
	0.116	161.0(4)
	0.118	161.9(4)
	0.120	162.8(4)
	0.122	163.5(4)

- Positive correlation between α_s and $m_t(m_t)$ **reduced once $t\bar{t}$ differential data included in the fit**

Fitted gluon PDF (work in progress)



- Significant reduction of the gluon PDF uncertainty once differential $t\bar{t}$ data are included
- The fitted gluon PDF is consistent with ABMP16
- Some tension between ATLAS and CMS is noticeable
- Work in progress on producing this fit in the xFitter framework

<https://gitlab.cern.ch/fitters/xfitter>, www.xfitter.org

Summary

- Measurements of $t\bar{t}$ production at LHC have very rich potential for phenomenology
 - ▶ provide information on m_t , α_S and gluon PDF
- We have extracted m_t^{pole} using most recent ATLAS and CMS Run-1,2,3 measurements of $t\bar{t}$ production with a few hundred MeV precision [arXiv:2311.05509]
 - ▶ however, pushing this to higher precision might require better understanding of the top quark mass definition used in MC to unfold the data to parton level (+ renormalon)
- Included most recent Run-2 measurements in ABMP fit [to appear arXiv:24XX.XXXXX]
 - ▶ the new data are consistent with ABMP16
 - ▶ significant ($\sim$ of factor 2) constraints on the gluon PDF at $x \gtrsim 0.1$
 - ▶ improved ($\sim$ of factor 2) determination of $m_t(m_t)$
 - ▶ reduced correlation between α_S and $m_t(m_t)$

Outlook

- Differential measurements of $t\bar{t}$ production seem to show some tension
 - ▶ ATLAS+CMS effort on combining differential $t\bar{t}$ data will be useful
 - ▶ more information on experimental correlations should be reported
- LHCb covers a complementary kinematic region sensitive to higher x , but no measurements unfolded to parton level (without cuts p_T , y of decay products) were done
 - ▶ would be useful to have such measurements from LHCb
- It would be very nice to have NNLO differential calculations for $t\bar{t}$ +jets
 - ▶ would allow direct constraints on α_S

BACKUP

Theoretical calculations with MATRIX framework

- Using “private” (custom) version of MATRIX provided by Javier Mazzitelli
- Interfaced to PineAPPL [Carrazza et al., JHEP 12 (2020) 108] to produce PDF interpolation grids which are further used in xFitter <https://gitlab.com/fitters/xfitter>
 - ▶ reproduce NNLO calculations using any PDF set and/or varied μ_r, μ_f in $\sim$ seconds
 - ▶ interface implemented privately and only for $t\bar{t}$ production
 - ▶ **no NNLO/NLO K-factors etc.**
- Further modifications to MATRIX to make possible runs with $\Delta\sigma_{t\bar{t}} < 1\%$
 - ▶ skip calculation of identical things (tailored for $t\bar{t}$)
 - ▶ adapted to DESY Bird Condor cluster and local multicore machines
 - ▶ technical fixes related to memory and disk space usage etc.
- We did 6 runs with $m_t^{\text{pole}} \in [165, 177.5]$ GeV with step of 2.5 GeV and $\Delta\sigma_{t\bar{t}} = 0.2\%$
 - ▶ $\approx 350\text{K} \times 6 \approx 2\text{M}$ CPU hours (200 years on single CPU, 3 months of real time)
 - ▶ for differential distributions, integration uncertainties in bins are $\lesssim 0.5\%$
- Differential distributions obtained with fixed $r_{\text{cut}} = 0.0015$ (q_T subtraction)
 - ▶ checked that extrapolation to $r_{\text{cut}} = 0$ for total cross section produces difference $< 1\%$ (see also S. Catani et al., JHEP 07 (2019) 100)
- $\mu_r = \mu_f = H_T/4$, $H_T = \sqrt{m_t^2 + p_T^2(t)} + \sqrt{m_t^2 + p_T^2(\bar{t})}$, varied up and down by factor 2 with $0.5 \leq \mu_r/\mu_f \leq 2$ (7-point method)
- For PDF and top quark mass fit, m_t^{pole} is converted to $m_t(m_t)$, as customary in ABMP16

Data in ABMP16 fit [Alekhin et al., arXiv:1701:05838] (1)

Experiment	Process	Reference	NDP	χ^2
DIS				
HERA I+II	$e^\pm p \rightarrow e^\pm X$ $e^\pm p \rightarrow \nu^\pm X$	[61]	1168	1510
BCDMS	$\mu^+ p \rightarrow \mu^+ X$	[61]	351	411
NMC	$\mu^+ p \rightarrow \mu^+ X$	[60]	245	343
SLAC-49a	$e^- p \rightarrow e^- X$	[54, 62]	38	59
SLAC-49b	$e^- p \rightarrow e^- X$	[54, 62]	154	171
SLAC-87	$e^- p \rightarrow e^- X$	[54, 62]	109	103
SLAC-89b	$e^- p \rightarrow e^- X$	[56, 62]	90	79

DIS heavy-quark production

HERA I+II	$e^\pm p \rightarrow e^\pm cX$	[63]	52	62
H1	$e^\pm p \rightarrow e^\pm bX$	[15]	12	5
ZEUS	$e^\pm p \rightarrow e^\pm bX$	[16]	17	16
CCFR	$\nu p \rightarrow \mu^\pm cX$	[64]	89	62
CHORUS	$\nu p \rightarrow \mu^\pm cX$	[18]	6	7.6
NOMAD	$\nu p \rightarrow \mu^\pm cX$	[17]	48	59
NuTeV	$\nu p \rightarrow \mu^\pm cX$	[64]	89	49



DY

FNAL-605	$pCu \rightarrow \mu^+ \mu^- X$	[67]	119	165
FNAL-866	$pp \rightarrow \mu^+ \mu^- X$ $pD \rightarrow \mu^+ \mu^- X$	[68]	39	53

Top-quark production

ATLAS, CMS	$pp \rightarrow t\bar{t}X$	[27, 32]	10	2.3
CDF&DØ	$\bar{p}p \rightarrow t\bar{t}X$ $\bar{p}p \rightarrow t\bar{t}X$	[53]	2	1.1
ATLAS, CMS	$pp \rightarrow t\bar{t}X$	[33, 52]	23	13
CDF&DØ	$\bar{p}p \rightarrow t\bar{t}X$	[53]	1	0.2

$t\bar{t}$ and single top LHC data have been updated to latest LHCTOPWG:



Data in ABMP16 fit [Alekhin et al., arXiv:1701:05838] (2)



Experiment		ATLAS		CMS		DØ		LHCb	
$\sqrt{s}$ (TeV)		7	13	7	8	1.96		7	8
Final states		$W^+ \rightarrow l^+ \nu$ $W^- \rightarrow l^- \nu$ $Z \rightarrow l^+ l^-$	$W^+ \rightarrow l^+ \nu$ $W^- \rightarrow l^- \nu$ $Z \rightarrow l^+ l^-$	$W^+ \rightarrow \mu^+ \nu$ $W^- \rightarrow \mu^- \nu$ (asym)	$W^+ \rightarrow \mu^+ \nu$ $W^- \rightarrow \mu^- \nu$	$W^+ \rightarrow \mu^+ \nu$ $W^- \rightarrow \mu^- \nu$ (asym)	$W^+ \rightarrow e^+ \nu$ $W^- \rightarrow e^- \nu$ (asym)	$W^+ \rightarrow \mu^+ \nu$ $W^- \rightarrow \mu^- \nu$ $Z \rightarrow \mu^+ \mu^-$	$Z \rightarrow e^+ e^-$ $W^- \rightarrow \mu^- \nu$ $Z \rightarrow \mu^+ \mu^-$
Cut on the lepton P_T		$P_T^l > 20$ GeV	$P_T^e > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^e > 25$ GeV	$P_T^\mu > 20$ GeV	$P_T^e > 20$ GeV
Luminosity (1/fb)		0.035	0.081	4.7	18.8	7.3	9.7	1	2
Reference		[66]	[26]	[24]	[25]	[23]	[22]	[19]	[21]
NDP		30	6	11	22	10	13	31	17
χ^2	present analysis ^a	31.0	9.2	22.4	16.5	17.6	19.0	45.1	21.7
	CJ15 [6]	—	—	—	—	20	29	—	—
	CT14 [7]	42	—	— ^b	—	—	34.7	—	—
	JR14 [8]	—	—	—	—	—	—	—	—
	HERAFitter [197]	—	—	—	—	13	19	—	—
	MMHT14 [9]	39	—	—	—	21	—	—	—
	NNPDF3.0 [10]	35.4	—	18.9	—	—	—	—	—

^a The ABM12 [1] analysis has used older data sets from CMS and LHCb.


^b For the statistically less significant data with the cut of $P_T^\mu > 35$ GeV the value of $\chi^2 = 12.1$ was obtained.

TABLE VI: Compilation of precise data on W - and Z -boson production in pp and $p\bar{p}$ collisions and the χ^2 values obtained for these data sets in different PDF analyses using their individual definitions of χ^2 . The NNLO fit results are quoted as a default, while the NLO values are given for the CJ15 [6] and HERAFitter [197] PDFs. Missing table entries indicate that the respective data sets have not been used in the analysis.

$\sigma(t\bar{t})$ and single t in ABMP16tt PDF fit




$\sigma(t\bar{t})$ vs NNLO predictions using different PDFs




- Fixed $m_t^{\text{pole}} = 172.5$ GeV, $\mu_r = \mu_f = H_T/4$
- Reported χ^2 values with (and without) PDF uncertainties
- All PDF sets describe data reasonably well (depends on m_t^{pole} , α_S)
- Sensitivity to PDFs reduces with increasing $\sqrt{s}$ (lower x probed)

$\sigma(t\bar{t})$ vs NNLO predictions using different m_t^{pole}



- ABMP16, fixed $\mu_r = \mu_f = H_T/4$
- Change of m_t^{pole} by 1 GeV $\rightarrow$ change of $\sigma(t\bar{t})$ by $\approx 3\%$
- Preferable $m_t^{\text{pole}} \sim 170\text{--}172.5 \text{ GeV}$ (depends on PDF and α_S)

$\sigma(t\bar{t})$ vs NNLO predictions with scale variations




• ABMP16, fixed $m_t^{pole} = 172.5$ GeV

• Scale variations $^{+3\%}_{-5\%}$:

- ▶ larger than data uncertainty (best data uncertainty $\pm 1.9\%$)
- ▶ limit precision of m_t^{pole} extraction to 1 GeV
- ▶ can be reduced by using e.g. $\overline{MS}$ mass $m_t(m_t)$ EPJ C74 (2014) 3167, JHEP04 (2021) 043

Extraction of m_t^{pole} : summary




- Extracted m_t^{pole} values with precision ± 0.3 GeV are consistent with PDG value 172.5 ± 0.7 GeV
 - ▶ data uncertainty ~ 0.2 GeV
 - ▶ PDF uncertainty ~ 0.1 GeV
 - ▶ NNLO scale uncertainty ~ 0.2 GeV
- Significant dependence on PDFs (~ 0.5 GeV):
 - ▶ different m_t^{pole} used in different PDFs
 - ▶ PDFs, m_t^{pole} , α_s should be determined simultaneously
- For CMS 1904.05237, NNLO results are consistent with published results obtained at NLO
 - ▶ good convergence of perturbative series
- Larger sensitivity comes from differential data
 - ▶ 2D differential x-sections in $M(t\bar{t})$, $y(t\bar{t})$ constrain m_t^{pole} , PDFs and (indirectly) α_s
 - ▶ ideally, 3D cross section in $M(t\bar{t})$, $y(t\bar{t})$ and number of extra jets constrain α_s directly, but NNLO not yet available for $t\bar{t}$ +jets
- Possible effects from Coulomb and soft-gluon resummation near the $t\bar{t}$ production threshold are neglected: might be ~ 1 GeV?


[CMS Coll. EPJ C80 (2020) 658; Kiyo, Kuhn, Moch, Steinhauser, Uwer EPJ C60 (2009) 375; Mäkelä, Hoang, Lipka, Moch 2301.03546]

Extraction of m_t^{pole} : differential Run 1, Run 2



Run 1 differential




Run 2 differential




Extraction of m_t^{pole} : dilepton vs semileptonic, ATLAS vs CMS




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions using different PDFs




Data vs NNLO predictions using different m_t^{pole}




Data vs NNLO predictions for CMS 2402.08486




r cut variation in bins of TOP-20-001



PineAPPL vs MATRIX in bins of TOP-20-001 [ABMP16]




PineAPpl vs MATRIX in bins of TOP-20-001 [CT18]




● grids were produced with ABMP16


MATRIX vs HighTEA



CMS TOP-18-004 checks



DESY 2018 summer school, L. Materne, bachelor thesis
 “Differential Top-Pair Production Cross Section with the CMS
 Detector - Optimization of Measurement Information”,
 Karlsruher Institut für Technologie (KIT), Bachelorarbeit,
 2018 [ETP-Bachelor-KA/2018-11]



m_t dependence of measured cross sections [CMS TOP-18-004]

