


Towards a Definitive Identification of the PTA Signal Origin


Andrea Mitridate

Towards a realistic forecast detection of primordial gravitational wave backgrounds, Valencia | Dec 13, 2024



NANOGrav
Physics Frontiers Center


TIMING RESIDUALS



**Pulses Recorded
by Radio Telescope**

GW SIGNALS

$$\frac{\delta P_a(t)}{P_a} = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega} \cdot \hat{p}_a)} \left[h_{ij}(t, \vec{0}) - h_{ij}(t - L_a, L_a \hat{p}_a) \right]$$



GW SIGNALS


$$\frac{\delta P_a(t)}{P_a} = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega} \cdot \hat{p}_a)} \left[h_{ij}(t, \vec{0}) - h_{ij}(t - L_a, L_a \hat{p}_a) \right]$$


Earth term

pulsar term

geometric response

$$\delta t_a(t) = \int^t dt' \frac{\delta P_a(t')}{P_a}$$





CONTINUOUS WAVE

$$h_{ij}(t, \vec{x}) = \sum_A h^A(t - \hat{\Omega} \cdot \vec{x}) e_{ij}^A(\hat{\Omega})$$

assuming:

- constant orbital frequency, ω
- plane of the orbit is $\perp$ to the line of sight

$$h^+(x) = h_0 \cos(\omega x + \phi) \quad h^\times(x) = h_0 \sin(\omega x + \phi)$$

CONTINUOUS WAVE



$$\delta t \propto F_A(h_A^e - h_A^p)$$

$$h_{ij}(t, \vec{x}) = \sum_A h^A(t - \hat{\Omega} \cdot \vec{x}) e_{ij}^A(\hat{\Omega})$$

assuming:

- constant orbital frequency, ω
- plane of the orbit is $\perp$ to the line of sight

$$h^+(x) = h_0 \cos(\omega x + \phi) \quad h^\times(x) = h_0 \sin(\omega x + \phi)$$





GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \tilde{h}_A(f, \hat{\Omega}) e_{ij}^A(\hat{\Omega}) e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}$$

GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \underbrace{\tilde{h}_A(f, \hat{\Omega})}_{\text{"Fourier" components}} \underbrace{e_{ij}^A(\hat{\Omega})}_{\text{polarization tensors}} \underbrace{e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}}_{\text{plane waves}}$$

GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \underbrace{\tilde{h}_A(f, \hat{\Omega})}_{\text{"Fourier" components}} \underbrace{e_{ij}^A(\hat{\Omega})}_{\text{polarization tensors}} \underbrace{e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}}_{\text{plane waves}}$$

the complex functions $\tilde{h}^A(f, \hat{\Omega})$ are treated as random variables

GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \underbrace{\tilde{h}_A(f, \hat{\Omega})}_{\text{"Fourier" components}} \underbrace{e_{ij}^A(\hat{\Omega})}_{\text{polarization tensors}} \underbrace{e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}}_{\text{plane waves}}$$

the complex functions $\tilde{h}^A(f, \hat{\Omega})$ are treated as random variables

$$\langle \tilde{h}_A(f, \hat{\Omega}) \rangle = 0$$

GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \underbrace{\tilde{h}_A(f, \hat{\Omega})}_{\text{"Fourier" components}} \underbrace{e_{ij}^A(\hat{\Omega})}_{\text{polarization tensors}} \underbrace{e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}}_{\text{plane waves}}$$

the complex functions $\tilde{h}^A(f, \hat{\Omega})$ are treated as random variables

$$\langle \tilde{h}_A(f, \hat{\Omega}) \rangle = 0$$

$$\langle \tilde{h}_A(f, \hat{\Omega})^* \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(\hat{\Omega}, \hat{\Omega}') \delta(f - f') \underbrace{H(f)}_{\text{GWB power spectrum}}$$

GW BACKGROUND

$$h_{ij}(t, \vec{x}) = \sum_A \int df \int d\hat{\Omega} \underbrace{\tilde{h}_A(f, \hat{\Omega})}_{\text{"Fourier" components}} \underbrace{e_{ij}^A(\hat{\Omega})}_{\text{polarization tensors}} \underbrace{e^{-2\pi i f(t - \hat{\Omega} \cdot \vec{x})}}_{\text{plane waves}}$$

the complex functions $\tilde{h}^A(f, \hat{\Omega})$ are treated as random variables

$$\langle \tilde{h}_A(f, \hat{\Omega}) \rangle = 0$$

$$\langle \delta t_a(t) \rangle = 0$$



$$\langle \tilde{h}_A(f, \hat{\Omega})^* \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(\hat{\Omega}, \hat{\Omega}') \delta(f - f') \underbrace{H(f)}_{\text{GWB power spectrum}}$$

$$\langle \delta t_a(t_i) \delta t_b(t_j) \rangle \propto \Gamma_{ab} \int df H(f) e^{2\pi i f(t_i - t_j)}$$

GW BACKGROUND

$$\langle \delta t_a(t_i) \delta t_b(t_j) \rangle \propto \Gamma_{ab} \int df \, H(f) \, e^{2\pi i f(t_i - t_j)}$$

GW BACKGROUND

$$\langle \delta t_a(t_i) \delta t_b(t_j) \rangle \propto \Gamma_{ab} \int df H(f) e^{2\pi i f(t_i - t_j)}$$

the signal is time-correlated



GW BACKGROUND

$$\langle \delta t_a(t_i) \delta t_b(t_j) \rangle \propto \Gamma_{ab} \int df H(f) e^{2\pi i f(t_i - t_j)}$$

the signal is time-correlated



the **power spectrum** of the GWB-induced signal is **common across pulsars**

GW BACKGROUND


$$\langle \delta t_a(t_i) \delta t_b(t_j) \rangle \propto \Gamma_{ab} \int df H(f) e^{2\pi i f(t_i - t_j)}$$

the signal is **correlated among pulsars**

$$\Gamma_{ab} = \int d\hat{\Omega} \sum_A F_a^A(\hat{\Omega}) F_b^A(\hat{\Omega}) [1 + \delta_{ab}]$$

$$F_a^A(\hat{\Omega}) = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega} \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega})$$

HELLINGS & DOWNS CURVE





searching for a GWB means searching for **stochastic delays** in the TOAs that are **common across pulsars** and **spatially correlated**

EVIDENCE FOR GWB

NANOGrav 15-year

[Agazie et al. \[2306.16213\]](#)




we find evidence for Hellings & Downs correlation with a p -value of $5 \times 10^{-5} - 1.9 \times 10^{-4}$ (approx. $3.5 - 4\sigma$)

EVIDENCE FOR GWB


NANOGrav:

68 pulsars, 16yr of data
~3-4 σ significance




PPTA:

32 pulsars, 18yr of data
~2 σ significance




EPTA + InPTA:

25 pulsars, 24yr of data
~3 σ significance





CPTA:

57 pulsars, 3yr of data
~4.6 σ significance




SPECTRUM


NANOGrav





PPTA




Reardon et al. [2306.16215]




EPTA + InPTA



IPTA early data combination




Agazie et al. [2309.00693]



what is the source?


ASTRO OR COSMO?




VS




ASTRO OR COSMO?



VS




ASTRO OR COSMO?



VS




ASTRO OR COSMO?



VS



ASTRO OR COSMO?





VS




HOW TO TELL THE DIFFERENCE

what are the distinguishing features of astrophysical and cosmological signals?









HOW TO TELL THE DIFFERENCE


what are the distinguishing features of astrophysical and cosmological signals?






HOW TO TELL THE DIFFERENCE

what are the distinguishing features of astrophysical and cosmological signals?







a deep dive on anisotropies



THE BASIC IDEA

isotropic GWB


$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle \propto \delta(\hat{\Omega}, \hat{\Omega}')$$



THE BASIC IDEA

anisotropic GWB

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle \propto \delta(\hat{\Omega}, \hat{\Omega}') P(\hat{\Omega})$$



$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$

↑ pulsars correlations

↑ geometric response


← GWB power

$$F_{a,k}^A = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega}_k \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega}_k)$$

THE BASIC IDEA

anisotropic GWB

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle \propto \delta(\hat{\Omega}, \hat{\Omega}') P(\hat{\Omega})$$



$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$

↑
pulsars
correlations

← GWB power


↑
geometric
response

$$F_{a,k}^A = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega}_k \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega}_k)$$

THE BASIC IDEA

anisotropic GWB

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle \propto \delta(\hat{\Omega}, \hat{\Omega}') P(\hat{\Omega})$$



$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$

← GWB power

↑
pulsars
correlations


↑
geometric
response

$$F_{a,k}^A = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega}_k \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega}_k)$$

THE BASIC IDEA

anisotropic GWB

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle \propto \delta(\hat{\Omega}, \hat{\Omega}') P(\hat{\Omega})$$



$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$


↑
pulsars
correlations

← GWB power



↑
geometric
response

$$F_{a,k}^A = \frac{\hat{p}_a^i \hat{p}_a^j}{2(1 + \hat{\Omega}_k \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega}_k)$$


THE BASIC IDEA





P_k



P_k





$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$




THE BASIC IDEA





P_k



P_k



$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$



FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

$$\rho_{ab}$$

the noise

cross correlations uncertainties

$$\Sigma_{ab}$$

the sky map

measured cross correlations

$$P_k$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

$$\rho_{ab}$$

the noise

cross correlations uncertainties

$$\Sigma_{ab}$$

the sky map

measured cross correlations

$$P_k$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

$$\rho_{ab}$$

the noise

cross correlations uncertainties

$$\Sigma_{ab}$$

the sky map

measured cross correlations

$$P_k$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

$$\rho_{ab}$$

the noise

cross correlations uncertainties

$$\Sigma_{ab}$$

what we have

the sky map

measured cross correlations

$$P_k$$

what we want

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

ρ_{ab}

the noise

cross correlations uncertainties

Σ_{ab}

what we have

the sky map

measured cross correlations

P_k

what we want

likelihood to observe a set of correlations, ρ , given a sky map $\mathbf{P}$

$$p(\rho|\mathbf{P}) = \frac{\exp[-\frac{1}{2}(\rho - \mathbf{R}\mathbf{P})^T \mathbf{\Sigma}^{-1}(\rho - \mathbf{R}\mathbf{P})]}{\sqrt{\det(2\pi\mathbf{\Sigma})}}$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

ρ_{ab}

the noise

cross correlations uncertainties

Σ_{ab}

what we have

the sky map

measured cross correlations

P_k

what we want

likelihood to observe a set of correlations, ρ , given a sky map $\mathbf{P}$

$$p(\rho|\mathbf{P}) = \frac{\exp[-\frac{1}{2}(\rho - \mathbf{R}\mathbf{P})^T \mathbf{\Sigma}^{-1}(\rho - \mathbf{R}\mathbf{P})]}{\sqrt{\det(2\pi\mathbf{\Sigma})}}$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

ρ_{ab}

the noise

cross correlations uncertainties

Σ_{ab}

what we have

the sky map

measured cross correlations

P_k

what we want

likelihood to observe a set of correlations, ρ , given a sky map $\mathbf{P}$

$$p(\rho|\mathbf{P}) = \frac{\exp[-\frac{1}{2}(\rho - \mathbf{R}\mathbf{P})^T \mathbf{\Sigma}^{-1}(\rho - \mathbf{R}\mathbf{P})]}{\sqrt{\det(2\pi\mathbf{\Sigma})}}$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

ρ_{ab}

the noise

cross correlations uncertainties

Σ_{ab}

what we have

the sky map

measured cross correlations

P_k

what we want

likelihood to observe a set of correlations, ρ , given a sky map $\mathbf{P}$

$$p(\rho|\mathbf{P}) = \frac{\exp[-\frac{1}{2}(\rho - \mathbf{R}\mathbf{P})^T \mathbf{\Sigma}^{-1}(\rho - \mathbf{R}\mathbf{P})]}{\sqrt{\det(2\pi\mathbf{\Sigma})}}$$

$$\mathbf{R} \propto F_{a,k}^A F_{b,k}^A$$

FREQUENTIST SEARCHES FOR ANISOTROPIES

three ingredients

the data

measured cross correlations

ρ_{ab}

the noise

cross correlations uncertainties

Σ_{ab}

what we have

the sky map


measured cross correlations

P_k

what we want


likelihood to observe a set of correlations, ρ , given a sky map $\mathbf{P}$

$$p(\rho|\mathbf{P}) = \frac{\exp[-\frac{1}{2}(\rho - \mathbf{R}\mathbf{P})^T \mathbf{\Sigma}^{-1}(\rho - \mathbf{R}\mathbf{P})]}{\sqrt{\det(2\pi\mathbf{\Sigma})}}$$



given $\{\rho, \mathbf{\Sigma}\}$ maximize p with respect to $\mathbf{P}$

FREQUENTIST SEARCHES FOR ANISOTROPIES



DETECTION STATISTIC

given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

DETECTION STATISTIC

given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

$$\text{SNR} \equiv \sqrt{2 \ln \left[\frac{p(\rho | \hat{\mathbf{P}})}{p(\rho | \mathbf{1})} \right]}$$

DETECTION STATISTIC

given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

$$\text{SNR} \equiv \sqrt{2 \ln \left[\frac{p(\rho | \hat{\mathbf{P}})}{p(\rho | \mathbf{1})} \right]}$$


how do we assess the statistical significance of a measured SNR? we need a null distribution!

DETECTION STATISTIC

given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

$$\text{SNR} \equiv \sqrt{2 \ln \left[\frac{p(\rho | \hat{\mathbf{P}})}{p(\rho | \mathbf{1})} \right]}$$

how do we assess the statistical significance of a measured SNR? we need a null distribution!




DETECTION STATISTIC

given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

$$\text{SNR} \equiv \sqrt{2 \ln \left[\frac{p(\rho | \hat{\mathbf{P}})}{p(\rho | \mathbf{1})} \right]}$$

how do we assess the statistical significance of a measured SNR? we need a null distribution!




DETECTION STATISTIC




given a reconstructed sky map, $\hat{\mathbf{P}}$, how do we quantify the evidence for anisotropies? we need a detection statistic!

$$\text{SNR} \equiv \sqrt{2 \ln \left[\frac{p(\rho | \hat{\mathbf{P}})}{p(\rho | \mathbf{1})} \right]}$$




how do we assess the statistical significance of a measured SNR? we need a null distribution!



GENERATING NULL DISTRIBUTIONS



GENERATING NULL DISTRIBUTIONS




one realization of the null hypothesis

$$\rho_{ab} = \text{HD}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$






"true value"



noise

GENERATING NULL DISTRIBUTIONS



one realization of the null hypothesis

$$\rho_{ab} = \text{HD}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$




↑
"true value"

↑
noise



reconstruct sky map
and get SNR

GENERATING NULL DISTRIBUTIONS



one realization of the null hypothesis



$$\rho_{ab} = \text{HD}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$

↑
"true value"


↑
noise




reconstruct sky map
and get SNR



WHY IS LIFE SO HARD?!




WHY IS LIFE SO HARD?!



WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$




WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables




WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can
be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$


one **amplitude** and **phase** for
each, **pixel**, **frequency**, and
polarization

WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can
be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for
each, **pixel**, **frequency**, and
polarization


given a set of $\{h_{\mathbf{k},i}^A, \phi_{\mathbf{k},i}^A\}$, the cross correlation coefficients can be uniquely derived

WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A}_{\text{random variables}} e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can
be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for
each, **pixel**, **frequency**, and
polarization

given a set of $\{h_{\mathbf{k},i}^A, \phi_{\mathbf{k},i}^A\}$, the cross correlation coefficients can be uniquely derived


ugly equation incoming

WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can
be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for
each, **pixel**, **frequency**, and
polarization

given a set of $\{h_{\mathbf{k},i}^A, \phi_{\mathbf{k},i}^A\}$, the cross correlation coefficients can be uniquely derived


$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} R_{akj}^{A*} R_{bk'j'}^{A'} h_{kj}^A h_{k'j'}^{A'} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$

WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**

given a set of $\{h_{\mathbf{k},i}^A, \phi_{\mathbf{k},i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} R_{akj}^{A*} R_{bk'j'}^{A'} h_{kj}^A h_{k'j'}^{A'} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$


sum over frequencies,
pixels, and polarizations

WHY IS LIFE SO HARD?!


each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A}_{\text{random variables}} e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can be thought as a collection of



given a set of $\{h_{k,i}^A, \phi_{k,i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} \boxed{R_{akj}^{A*} R_{bk'j'}^{A'}} h_{kj}^A h_{k'j'}^{A'} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$

response functions for
pulsars a and b

$$R_a^A(f, \hat{\Omega}) \equiv F_a^A(\hat{\Omega}) \left[1 - e^{2\pi i f L_a (1 + \hat{p}_a \cdot \hat{\Omega})} \right]$$


$$F_a^A(\hat{\Omega}) \equiv \frac{p_a^i p_a^j}{2(1 + \hat{\Omega} \cdot \hat{p}_a)} e_{ij}^A(\hat{\Omega})$$

WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \boxed{h_A} e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \boxed{\phi})}$$

random variables



one realization of a GWB can be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**

given a set of $\{h_{k,i}^A, \phi_{k,i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} R_{akj}^{A*} R_{bk'j'}^{A'} \boxed{h_{kj}^A h_{k'j'}^{A'}} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$


pixel amplitudes

WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}}_{\text{random variables}}$$

random variables



one realization of a GWB can be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**

given a set of $\{h_{k,i}^A, \phi_{k,i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} R_{akj}^{A*} R_{bk'j'}^{A'} h_{kj}^A h_{k'j'}^{A'} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$


interference between different pixels

WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}}_{\text{random variables}}$$

random variables



one realization of a GWB can be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**

given a set of $\{h_{\mathbf{k},i}^A, \phi_{\mathbf{k},i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\rho_{ab} = \Delta f^2 \Delta \hat{\Omega}^2 \sum_{jj'} \sum_{kk'} \sum_{AA'} R_{akj}^{A*} R_{bk'j'}^{A'} h_{kj}^A h_{k'j'}^{A'} \exp(i\Delta\phi_{kk',jj'}^{AA'}) \text{sinc}(\pi(j - j')) + \text{c.c.}$$


interference between different frequencies

WHY IS LIFE SO HARD?!


each of this pixel can be thought as emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A}_{\text{random variables}} e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can be thought as a collection of



given a set of $\{h_{k,i}^A, \phi_{k,i}^A\}$, the cross correlation coefficients can be uniquely derived

$$\langle \rho_{ab} \rangle \propto \sum_A \sum_k F_{a,k}^A F_{b,k}^A P_k$$


if we average over many realizations of the GWB this reduces to the usual expression

WHY IS LIFE SO HARD?!

each of this pixel can be thought as
emitting a GW plane wave

$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}}_{\text{random variables}}$$


random variables



one realization of a GWB can
be thought as a collection of

$$h_{\mathbf{k},i}^A \quad \phi_{\mathbf{k},i}^A$$

one **amplitude** and **phase** for
each, **pixel**, **frequency**, and
polarization




each realization of an isotropic GWB
will produce cross correlation that
fluctuate around the HD value

WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave



$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A^A e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}}_{\text{random variables}}$$

random variables



one realization of a GWB can be thought as a collection of

$h_{\mathbf{k},i}^A$ $\phi_{\mathbf{k},i}^A$
 one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**




WHY IS LIFE SO HARD?!

each of this pixel can be thought as emitting a GW plane wave


$$h_{ij}(t, \vec{x}) = \sum_A \underbrace{h_A}_{\text{random variables}} e_{ij}^A(\mathbf{k}) e^{i2\pi f(t - \mathbf{k} \cdot \mathbf{x} + \phi)}$$

random variables



one realization of a GWB can be thought as a collection of

$h_{\mathbf{k},i}^A$
 $\phi_{\mathbf{k},i}^A$
 one **amplitude** and **phase** for each, **pixel**, **frequency**, and **polarization**



NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$

take one realization of the
null hypothesis

$$\{h_{k,i}^A, \phi_{k,i}^A\}$$


NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$

take one realization of the
null hypothesis

$$\{h_{k,i}^A, \phi_{k,i}^A\}$$



compute induced cross
correlations

$$\hat{\rho}_{ab} = \text{ugly_equation}(h_{k,i}^A, \phi_{k,i}^A)$$


NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$


take one realization of the
null hypothesis

$$\{h_{k,i}^A, \phi_{k,i}^A\}$$



compute induced cross
correlations

$$\hat{\rho}_{ab} = \text{ugly_equation}(h_{k,i}^A, \phi_{k,i}^A)$$



add noise

$$\rho_{ab} = \hat{\rho}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$


NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$


take one realization of the
null hypothesis

$$\{h_{k,i}^A, \phi_{k,i}^A\}$$




compute induced cross
correlations

$$\hat{\rho}_{ab} = \text{ugly_equation}(h_{k,i}^A, \phi_{k,i}^A)$$



add noise

$$\rho_{ab} = \hat{\rho}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$




run anisotropy search on
these mock correlations and
get an reconstructed sky

NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$




NULL DISTRIBUTIONS v2.0

given an isotropic GWB as null hypothesis

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_A(f', \hat{\Omega}') \rangle = \delta_{AA'} \delta(f - f') \delta(\hat{\Omega}, \hat{\Omega}') H(f)$$


take one realization of the
null hypothesis

$$\{h_{k,i}^A, \phi_{k,i}^A\}$$




compute induced cross
correlations

$$\hat{\rho}_{ab} = \text{ugly_equation}(h_{k,i}^A, \phi_{k,i}^A)$$




add noise



$$\rho_{ab} = \hat{\rho}_{ab} + \mathcal{N}(0, \Sigma_{ab})$$




run anisotropy search on
these mock correlations and
get a reconstructed sky



compute anisotropic SNR




NULL DISTRIBUTIONS v2.0




Konstandin, Lemke, AM, Perboni *"The impact of cosmic variance on PTAs anisotropy searches"*, [2498.07741]

NULL DISTRIBUTIONS v2.0



Konstandin, Lemke, AM, Perboni *"The impact of cosmic variance on PTAs anisotropy searches"*, [2498.07741]


SO, HAVE WE DETECTED ANISOTROPIES?!



for the reconstructed GWB map in the NG15 data

$$\text{SNR} \sim 4 \Rightarrow p \sim 0.05$$

SO, HAVE WE DETECTED ANISOTROPIES?!




for the reconstructed GWB map in the NG15 data

$$\text{SNR} \sim 4 \Rightarrow p \sim 0.5 \text{ } \times \text{ } 0.7$$


INTERFERENCE, INTERFERENCE EVERYWHERE

how astrophysical inference analyses are performed




INTERFERENCE, INTERFERENCE EVERYWHERE

how astrophysical inference analyses are performed




INTERFERENCE, INTERFERENCE EVERYWHERE

how astrophysical inference analyses are performed




INTERFERENCE, INTERFERENCE EVERYWHERE

how astrophysical inference analyses are performed




INTERFERENCE, INTERFERENCE EVERYWHERE

how astrophysical inference analyses are performed




INTERFERENCE, INTERFERENCE EVERYWHERE

$$\left\{ \tilde{h}_i^A, f_i, \Omega_i \right\}$$




expected
 $\Phi_j \propto \langle \tilde{\delta} t_j \tilde{\delta} t_j \rangle$




$$\Phi_j \propto \sum_{i \in \text{bin } j} |\tilde{h}_i|^2$$

INTERFERENCE, INTERFERENCE EVERYWHERE

$$\left\{ \tilde{h}_i^A, f_i, \Omega_i \right\}$$



expected
 $\Phi_j \propto \langle \tilde{\delta} t_j \tilde{\delta} t_j \rangle$




$$\Phi_j \propto \sum_{ii'} \tilde{h}_i \tilde{h}_{i'} w_{ji} w_{ji'}$$

INTERFERENCE, INTERFERENCE EVERYWHERE


how astrophysical inference analyses are performed

Xue et al. [2409.19516]



The background of the slide is black, featuring numerous abstract, colorful circular patterns. These patterns are composed of multiple overlapping lines in shades of teal, orange, and purple, creating a sense of depth and movement. They are scattered across the entire frame, with some appearing larger and more prominent than others.

strong evidence for a GWB in the nHz band



strong evidence for a GWB in the nHz band

cosmology or astrophysics?



strong evidence for a GWB in the nHz band

cosmology or astrophysics?

CW and anisotropies will help us discriminating

strong evidence for a GWB in the nHz band

cosmology or astrophysics?

CW and anisotropies will help us discriminating


we need to develop tools to characterize the GWB

backup

HOW TO SAMPLE A GWB

$$\langle \tilde{h}_{kj}^{A*} \tilde{h}_{k'j'}^{A'} \rangle = \delta_{AA'} \delta_{kk'} \delta_{jj'} \frac{H_j}{\Delta \hat{\Omega} \Delta f}$$

$$\langle \tilde{h}_{kj}^A \tilde{h}_{k'j'}^{A'} \rangle = \langle \tilde{h}_{kj}^{A*} \tilde{h}_{k'j'}^{A'*} \rangle = 0$$





$$\tilde{h}_{kj}^A \equiv h_{kj}^A e^{i\phi_{kj}^A}$$

$$p(h) = \frac{h}{2\pi\sigma_j^2} e^{-h^2/2\sigma_j^2}$$


$$\sigma_j^2 = H_j / (2\Delta \hat{\Omega} \Delta f)$$


$$p(\phi) = \text{Uniform}[0, 2\pi]$$


RESULTS IN THE PIXEL BASIS




PULSAR TIMING




PULSAR TIMING




PULSAR TIMING



PULSAR TIMING




PULSAR TIMING



$$t_{\text{TOA}} = T_{\text{obs}} + \Delta\tau$$

PULSAR TIMING



$$t_{\text{TOA}} = T_{\text{obs}} + \Delta\tau$$

BUILDING A TIMING MODEL



$$t_e^{\text{psr}} = t_a^{\text{obs}} - \Delta_{\odot} - \Delta_{\text{IS}} - \Delta_{\text{B}}$$

the goal is to relate the arrival time in Terrestrial Time (TT) to the emission time in the pulsar reference frame

BUILDING A TIMING MODEL

$$t_e^{\text{psr}} = t_a^{\text{obs}} - \boxed{\Delta_{\odot}} - \Delta_{\text{IS}} - \Delta_{\text{B}}$$

first step transform from the from the TT time to the SSB time



$$\begin{aligned} \Delta_{\odot} = & \Delta_{\text{A}} \\ & + \\ & \Delta_{\text{R}\odot} \\ & + \\ & \Delta_{\text{p}} \\ & + \\ & \Delta_{\text{D}\odot} \\ & + \\ & \Delta_{\text{S}\odot} \\ & + \\ & \Delta_{\text{E}\odot} \end{aligned}$$

atmospheric delay: group velocity of radio waves in the atmosphere differs from the vacuum speed of light

Roemer delay: difference in the arrival time of the pulse at the observatory and at the SSB

Parallax term: delay due to the curvature of spherical wavefront

dispersion term: due to radio waves propagation in interplanetary medium

Shapiro delay: due to propagation of radio pulses through potential wells of solar systems bodies

Einstein delay: difference in the arrival time due to acceleration and potential of the Earth

BUILDING A TIMING MODEL


$$t_e^{\text{psr}} = t_a^{\text{obs}} - \Delta_{\odot} - \Delta_{\text{IS}} - \Delta_{\text{B}}$$

second step transform from the from SSB to the BB time

$$\Delta_{\text{IS}} = \Delta_{\text{VP}} \quad \text{vacuum propagation delay: vacuum light travel time}$$

$$\Delta_{\text{ISD}} \quad \text{dispersion term: due to propagation through the interstellar medium}$$

$$\Delta_{\text{ES}} \quad \text{Einstein delay: due to the secular motion of the SSB}$$




BUILDING A TIMING MODEL

$$t_e^{\text{psr}} = t_a^{\text{obs}} - \Delta_{\odot} - \Delta_{\text{IS}} - \Delta_{\text{B}}$$

last step transform from the from BB to the pulsar time

$$\Delta_{\text{B}} = \Delta_{\text{RB}}$$

Roemer delay: delay due to the orbital motion of the pulsar




$$\Delta_{\text{EB}}$$

Einstein delay: difference in the emission time from the pulsar inertial reference frame and the binary barycenter

$$\Delta_{\text{SB}}$$


Shapiro delay: effect of the gravitational potential of the binary companion



BUILDING A TIMING MODEL

$$t_e^{\text{psr}} = t_a^{\text{obs}} - \Delta_{\odot} - \Delta_{\text{IS}} - \Delta_{\text{B}}$$

[arXiv: 0603381](#)




Correction	Typical value/range
Observatory clock to TT	1 μ s
Hydrostatic tropospheric delay	10 ns
Zenith wet delay	1.5 ns
IAU precession/nutation	$\sim$ 5 ns
Polar motion	60 ns
Δ UT1	1 μ s
Einstein delay	1.6 ms
Roemer delay	500 s
Shapiro delay due to Sun	112 μ s
Shapiro delay due to Venus	0.5 ns
Shapiro delay due to Jupiter	180 ns
Shapiro delay due to Saturn	58 ns
Shapiro delay due to Uranus	10 ns
Shapiro delay due to Neptune	12 ns
Second order Solar Shapiro delay	9 ns
Interplanetary medium dispersion delay	100 ns ^b
Interstellar medium dispersion delay	$\sim$ 1 s ^b

BUILDING A TIMING MODEL

$$\phi(t^{\text{psr}}) = \phi_0 + \nu_0(t^{\text{psr}} - t_0) + \frac{1}{2}\dot{\nu}_0(t^{\text{psr}} - t_0)^2$$



unknown quantities



BUILDING A TIMING MODEL

$$\phi(t^{\text{psr}}) = \phi_0 + \nu_0(t^{\text{psr}} - t_0) + \frac{1}{2}\dot{\nu}_0(t^{\text{psr}} - t_0)^2$$


unknown quantities




unknown timing models parameters are fitted to the TOAs

$$\chi^2 = \sum_i \left(\frac{\phi(t_{\text{e},i}^{\text{psr}}) - N_i}{\sigma_i} \right)^2$$

BUILDING A TIMING MODEL

$$\phi(t^{\text{psr}}) = \phi_0 + \nu_0(t^{\text{psr}} - t_0) + \frac{1}{2}\dot{\nu}_0(t^{\text{psr}} - t_0)^2$$


unknown quantities



unknown timing models parameters are fitted to the TOAs


$$\chi^2 = \sum_i \left(\frac{\phi(t_{\text{e},i}^{\text{psr}}) - N_i}{\sigma_i} \right)^2$$

closest integer

BUILDING A TIMING MODEL

$$\phi(t^{\text{psr}}) = \phi_0 + \nu_0(t^{\text{psr}} - t_0) + \frac{1}{2}\dot{\nu}_0(t^{\text{psr}} - t_0)^2$$

↑ ↑ ↑
unknown quantities



unknown timing models parameters are fitted to the TOAs

$$\chi^2 = \sum_i \left(\frac{\phi(t_{e,i}^{\text{psr}}) - N_i}{\sigma_i} \right)^2$$

closest integer

time of emission in the
pulsar reference frame

FACE-OFF

free parameters

$$h^2 \Omega_{\text{GW}} \propto \frac{A^2}{H_0^2} \left(\frac{f}{\text{yr}^{-1}} \right)^{5-\gamma} \text{yr}^{-2}$$

vs

free parameters

$$h^2 \Omega_{\text{GW}}(f; \Theta)$$

FACE-OFF

$$\mathcal{B} = \frac{\mathcal{Z}_{\text{NP}}}{\mathcal{Z}_{\text{BHB}}}$$

$$\mathcal{Z} = \int d\Theta \, P(\mathcal{D}|\Theta, \mathcal{H}) \times P(\Theta|\mathcal{H})$$

FACE-OFF

$$\mathcal{B} = \frac{\mathcal{Z}_{\text{NP}}}{\mathcal{Z}_{\text{BHB}}}$$

$$\mathcal{Z} = \int d\Theta \, P(\mathcal{D}|\Theta, \mathcal{H}) \times P(\Theta|\mathcal{H})$$



likelihood function

FACE-OFF

$$\mathcal{B} = \frac{\mathcal{Z}_{\text{NP}}}{\mathcal{Z}_{\text{BHB}}}$$


$$\mathcal{Z} = \int d\Theta \, P(\mathcal{D}|\Theta, \mathcal{H}) \times P(\Theta|\mathcal{H})$$

likelihood function

prior distributions

FACE-OFF

$$\mathcal{B} = \frac{\mathcal{Z}_{\text{NP}}}{\mathcal{Z}_{\text{BHB}}}$$



$$\mathcal{Z} = \int d\Theta \, P(\mathcal{D}|\Theta, \mathcal{H}) \times P(\Theta|\mathcal{H})$$

likelihood function

prior distributions