

# THE MILKY WAY'S FAINTEST SATELLITES

## THE CONTRIBUTION OF "ORPHAN" GALAXIES

**Dr Isabel Santos-Santos**

**Institute for Computational Cosmology, Durham University**

With Carlos Frenk, Julio Navarro, Shaun Cole, John Helly




Self-Interacting Dark Matter and Small-Scale Structure of the Universe - Valencia, June 2025



# Substructure in Cold Dark Matter

Aquarius MW-mass halo simulation [Springel+08]



| Name   | $m_p$<br>( $M_\odot$ ) | $\epsilon$<br>(pc) |
|--------|------------------------|--------------------|
| Aq-A-1 | $1.712 \times 10^3$    | 20.5               |


- 1 Well-defined **mass function**  
→ More small halos than large
- 2 Self-similar abundance of **substructure**
- 3 Universal, **"cuspy"** halo density profiles  
→ resilient to tidal effects
- 4 **Orbits** and accretion histories

[See e.g. Press&Schechter74; Bond+91; White&Rees78; White&Frenk91; NavarroFrenk&White97; Springel+08; Bullock&Boylan-Kolchin17].



# Substructure in Cold Dark Matter

Aquarius MW-mass halo simulation [Springel+08]



| Name   | $m_p$<br>( $M_\odot$ ) | $\epsilon$<br>(pc) |
|--------|------------------------|--------------------|
| Aq-A-1 | $1.712 \times 10^3$    | 20.5               |

- 1 Well-defined **mass function**  
→ More small halos than large
- 2 Self-similar abundance of **substructure**
- 3 Universal, "**cuspy**" halo density profiles  
→ resilient to tidal effects
- 4 **Orbits** and accretion histories

[See e.g. Press&Schechter74; Bond+91; White&Rees78; White&Frenk91; NavarroFrenk&White97; Springel+08; Bullock&Boylan-Kolchin17].



**\*\*Not up-to-date**



I.Santos-Santos (ICC, Durham)



# Substructure observed in the MW


\*\*Not up-to-date



Currently ~60 MW satellite candidates

Recent findings of extremely ultrafaint systems (e.g. Tuc3, UMa3/Unions1)

New upcoming deep surveys (e.g. Rubin LSST)



SDSS, DES, HSC, Gaia, DELVE, UNIONS  
e.g. Willman+05; Belokurov+06-10; Laevens+15; Bechtol+15;  
Drlica-Wagner15,21; Koposov+15; Homma+16; Mau+20;  
Cerny+21,23; Smith+22,23; Torrealba+19; Pace&Li19;  
McConnachie&Venn20.

Wikimedia Commons




# How to model the MW system in CDM?

**The unabridged satellite luminosity function of Milky Way-like galaxies in  $\Lambda$ CDM: the contribution of “orphan” satellites**

Isabel M.E. Santos-Santos,<sup>1</sup>★ Carlos S. Frenk,<sup>1</sup> Julio F. Navarro,<sup>2</sup> Shaun Cole,<sup>1</sup> John Helly<sup>1</sup>



## 1 Populate subhalos with galaxies



◆ A threshold for galaxy formation reflecting the minimum potential well depth needed for Hydrogen to cool  **$V_{\text{peak}} \sim 15\text{-}20 \text{ km/s}$** ;



*[Okamoto&Frenk19, Benitez-Llambay&Frenk20]*

◆ Only halos that reach this potential depth can form stars, rest remain dark.

◆ Implies number of MW satellites (of any luminosity) cannot be too high ( $< \sim 300$ )



### 2 Track substructure in time



- ◆ Many subhalos that should host satellites are lost due to numerical issues (artificially disrupt)
- ◆ In CDM, every subhalo survives tides, as protected by its cusp
- ◆ It is crucial to correct for artificial disruption and track “orphan” satellites

[see e.g. Peñarrubia+08,10; van den Bosch & Ogiya18; HAn+18; Green&van den Bosch 19; Jiang+21; Stucker+23; He+25; Errani+20,21,22,24]





# How to model the MW system in CDM?

The unabridged satellite luminosity function of Milky Way-like galaxies in  $\Lambda$ CDM: the contribution of “orphan” satellites

Isabel M.E. Santos-Santos,<sup>1\*</sup> Carlos S. Frenk,<sup>1</sup> Julio F. Navarro,<sup>2</sup> Shaun Cole,<sup>1</sup> John Helly<sup>1</sup>

## 2 Track substructure in time





- ◆ Many subhalos that should host satellites are lost due to numerical issues (artificially disrupt)
- ◆ In CDM, every subhalo survives tides, as protected by its cusp
- ◆ It is crucial to correct for artificial disruption and track “orphan” satellites

[see e.g. Peñarrubia+08,10; van den Bosch & Ogiya18; HAn+18; Green&van den Bosch 19; Jiang+21; Stucker+23; He+25; Errani+20,21,22,24]



## 2 The contribution of “orphan” satellites in MW halos




- ◆ “Orphans” dominate the inner regions of the MW
- ◆ Orphans make up **half** of the total satellite population, even in Aq-A-1



2

The contribution of “orphan” satellites in MW halos




|    | Particle mass<br>(Msun) | R(kpc) |
|----|-------------------------|--------|
| L1 | 1.7e+03                 | 30     |
| L2 | 1.4e+04                 | 69     |
| L4 | 4e+05                   | 102    |
| L5 | 3e+06                   | 150    |

- ◆ “Orphans” dominate the inner regions of the MW
- ◆ Orphans make up **half** of the total satellite population, even in Aq-A-1



2

The contribution of “orphan” satellites in MW halos



|    | Particle mass<br>(Msun) | R(kpc) |
|----|-------------------------|--------|
| L1 | 1.7e+03                 | 30     |
| L2 | 1.4e+04                 | 69     |
| L4 | 4e+05                   | 102    |
| L5 | 3e+06                   | 150    |

- ◆ “Orphans” dominate the inner regions of the MW
- ◆ Orphans make up **half** of the total satellite population, even in Aq-A-1



# How to model the MW system in CDM?

**GALFORM semi-analytical model**

[White&Frenk91, Kauffmann+93, Cole+2000, Lacey+2016]

**3** Model observed satellite properties

*Initial stellar mass and stellar size*

*Santos-Santos et al 2025; +in prep*





# How to model the MW system in CDM?

**GALFORM semi-analytical model**

[White&Frenk91, Kauffmann+93, Cole+2000, Lacey+2016]

## 3 Model observed satellite properties

*Initial stellar mass and stellar size*






# How to model the MW system in CDM?

## 3 Model observed satellite properties

*Tidal stripping on Dark Matter and stars*



[Errani+21,22]

**Subhalo tidal evolution in LCDM "Tidal Tracks"**


Gives DM mass loss



# How to model the MW system in CDM?


## 3 Model observed satellite properties

## Tidal stripping on Dark Matter and stars



**Subhalo tidal evolution in LCDM "Tidal Tracks"**

Gives DM mass loss



**Tidal evolution of the stellar component**


Gives change in galaxy luminosity and size



# How to model the MW system in CDM?


## 3 Model observed satellite properties

## Tidal stripping on Dark Matter and stars



**Subhalo tidal evolution in LCDM "Tidal Tracks"**


Gives DM mass loss



**Tidal evolution of the stellar component**


Gives change in galaxy luminosity and size






◆ Magnitudes have changed after stripping. Faint galaxies become much fainter





- ◆ Magnitudes have changed after stripping. Faint galaxies become much fainter
- ◆ The model predicts very well the bright end of the MW confirmed satellites Lum func






◆ Magnitudes have changed after stripping. Faint galaxies become much fainter

◆ The model predicts very well the bright end of the MW confirmed satellites Lum func

◆ Predicts a fairly large amount of faint satellites that are yet to be found ( $M_V > -3$  or  $M_{\text{star}} < 1000 M_{\text{sun}}$ )



# Predictions: MW satellite *luminosity function*




◆ Magnitudes have changed after stripping. Faint galaxies become much fainter

◆ The model predicts very well the bright end of the MW confirmed satellites Lum func

◆ Predicts a fairly large amount of faint satellites that are yet to be found ( $M_V > -3$  or  $M_{\text{star}} < 1000 M_{\text{sun}}$ )



# Predictions: MW satellite *luminosity function*




◆ Magnitudes have changed after stripping. Faint galaxies become much fainter

◆ The model predicts very well the bright end of the MW confirmed satellites Lum func

◆ Predicts a fairly large amount of faint satellites that are yet to be found ( $M_V > -3$  or  $M_{\text{star}} < 1000 M_{\text{sun}}$ )



# Predictions: MW satellite *luminosity function*



◆ Magnitudes have changed after stripping. Faint galaxies become much fainter


◆ The model predicts very well the bright end of the MW confirmed satellites Lum func

◆ Predicts a fairly large amount of faint satellites that are yet to be found ( $M_V > -3$  or  $M_{\text{star}} < 1000 M_{\text{sun}}$ )



# Newly discovered MW “faint ambiguous satellite candidates”

*Koposov et al 2015 (DES)*



- Found in imaging surveys. Just a few stars
- No line of sight velocities, or large errors. Unclear what is their nature (GCs or dwarfs?)

Koposov 2  
Koposov 1  
SEGUE 3  
Munoz 1  
Balbinot 1  
Kim 1  
Kim 2 (Indus 1)  
Draco 2  
Crater  
SMASH 1  
Kim 3  
DES J0111-1341  
DES J0225+0304  
Eridanus 3  
DES 1  
DES 3  
Pictor 1  
To 1  
Laevens 3  
DES 4  
PS1 1  
HSC 1  
DES 5  
BLISS 1  
Gaia 3  
DELVE 1  
DELVE 2  
YMCA-1  
DELVE 5  
DELVE 3  
DELVE 4  
DELVE 6  
UMa3/Unions 1  
...



# Newly discovered MW “faint ambiguous satellite candidates”




The faintest satellite known!

Smith et al 2024 (CFHT Unions Survey)  
See also Errani et al 2024

$M_{\text{star}} = 16^{+6}_{-5} M_{\odot}$

UMa3 / Unions1

$\sigma_v < 3.7^{+1.4}_{-1.0} \text{ km s}^{-1}$




- Found in imaging surveys. Just a few stars
- No line of sight velocities, or large errors. Unclear what is their nature (GCs or dwarfs?)

|                 |
|-----------------|
| Koposov 2       |
| Koposov 1       |
| SEGUE 3         |
| Munoz 1         |
| Balbinot 1      |
| Kim 1           |
| Kim 2 (Indus 1) |
| Draco 2         |
| Crater          |
| SMASH 1         |
| Kim 3           |
| DES J0111-1341  |
| DES J0225+0304  |
| Eridanus 3      |
| DES 1           |
| DES 3           |
| Pictor 1        |
| To 1            |
| Laevens 3       |
| DES 4           |
| PS1 1           |
| HSC 1           |
| DES 5           |
| BLISS 1         |
| Gaia 3          |
| DELVE 1         |
| DELVE 2         |
| YMCA-1          |
| DELVE 5         |
| DELVE 3         |
| DELVE 4         |
| DELVE 6         |
| UMa3/Unions 1   |
| ...             |




# Newly discovered MW “faint ambiguous satellite candidates”







# Newly discovered MW “faint ambiguous satellite candidates”





# Predictions: MW satellite *luminosities, sizes and distances*



Santos-Santos *in prep*





# Predictions: MW satellite *luminosities, sizes and distances*



Santos-Santos *in prep*





# Predictions: MW satellite *luminosities, sizes and distances*

Santos-Santos *in prep*







# Predictions: MW satellite *luminosities, sizes and distances*

Santos-Santos *in prep*

Data:[Smith+24]




$z=0$  Luminosities and sizes are consistent with data



No satellite with  $M_V < 5$  within 10 kpc of the MW centre




# Predictions: MW satellite *orbits* & *accretion times*



[see e.g. Battaglia+22, Pace, Erakal&Li22]





# Predictions: MW satellite *orbits* & *accretion times*



[see e.g. Battaglia+22, Pace, Erakal&Li22]









# Predictions: MW satellite *orbits* & *accretion times*

Santos-Santos *in prep*



[Santos-Santos+2025]



[see e.g. Battaglia+22, Pace, Erkal&Li22]



# Predictions: MW satellite *orbits* & *accretion times*



$$\bar{\rho}_{\text{host}}(r) \sim \bar{\rho}_{\text{sat}}$$

[see e.g. Battaglia+22, Pace, Erakal&Li22]



# Predictions: MW satellite *velocity dispersion*

[Errani+21 (Eq.8 & 9)]


**Truncated NFW profile.**

Depends on mass loss

$$\sigma = \sqrt{\frac{GM(< r_h)}{3r_h}}$$

$$M(< r_h) = (4/3)\pi r_h^3 \bar{\rho}(< r_h)$$

[see also Errani+24]





# Predictions: MW satellite *velocity dispersion*

[see also Errani+24]


[Errani+21 (Eq.8 & 9)]

**Truncated NFW profile.**

Depends on mass loss

$$\sigma = \sqrt{\frac{GM(< r_h)}{3r_h}}$$

$$M(< r_h) = (4/3)\pi r_h^3 \bar{\rho}(< r_h)$$





# Predictions: MW satellite *velocity dispersion*

[Errani+21 (Eq.8 & 9)]

**Truncated NFW profile.**


Depends on mass loss

$$\sigma = \sqrt{\frac{GM(< r_h)}{3r_h}}$$

$$M(< r_h) = (4/3)\pi r_h^3 \bar{\rho}(< r_h)$$

Errani+24 (Eq.2 self-gravitating)

$$\langle \sigma_{\text{los}}^2 \rangle = \frac{5}{96} \frac{GM_\star}{r_\star}$$



[see also Errani+24]



# Predictions: MW satellite *velocity dispersion*

[Errani+21 (Eq.8 & 9)]

**Truncated NFW profile.**


Depends on mass loss

$$\sigma = \sqrt{\frac{GM(< r_h)}{3r_h}}$$

$$M(< r_h) = (4/3)\pi r_h^3 \bar{\rho}(< r_h)$$

Errani+24 (Eq.2 self-gravitating)

$$\langle \sigma_{\text{los}}^2 \rangle = \frac{5}{96} \frac{GM_\star}{r_\star}$$



[see also Errani+24]


◆ Velocity dispersion measurement only way of discriminating conclusively if satellite is a GC or dwarf

◆ ...But it's difficult



## A) At first order:

- Heating from self-interactions leads to core formation.
- Subhalos affected by tides, prone to genuine disruption
- Host density profile also changes(!)



## B) With small fixed cross-sections ( $<0.1 \text{ cm}^2/\text{g}$ )

- Very small cores, improving survival of halos  $\rightarrow$  predictions similar to CDM

## C) With velocity-dependent large cross-sections ( $>10 \text{ cm}^2/\text{g}$ ) and "core-collapse":

- **C1)** If  $t_{\text{cc}} < T_{\text{infall}}$   $\rightarrow$  subhalo will resist tides  $\rightarrow$  prediction similar to CDM
- **C2)** If  $t_{\text{cc}} > T_{\text{infall}}$   $\rightarrow$  prediction depends on interplay between CC formation and tides



# Summary

- In **CDM, subhalos always survive**. Stars may be lost.
- Cosmological simulations suffer subhalo artificial disruption. It is crucial to correct for "orphans".
- LCDM models of the MW system match well the properties of satellite galaxies known so far. They also **predict the existence of numerous faint satellite galaxies at close distances**, with properties
  - $\sigma = 1\text{-}3 \text{ km/s}$
  - $M_V > -3$  ( $M_{\text{star}} < 10^3 M_{\text{sun}}$ )
  - $r_{\text{peri}} > 10 \text{ kpc}$ . ( $r_{\text{apo}} < \sim 100 \text{ kpc}$ )
- These properties seem to match those of "ambiguous" satellites identified recently with photometric surveys/Gaia
- Some of these predictions are unique to CDM, and may help to discriminate between dark matter models

Gracias!