


# The UHECR Mass Composition Measurements at the Pierre Auger Observatory

Berenika Čermáková for the Pierre Auger Collaboration



Credit: Pierre Auger Observatory









PIERRE  
AUGER  
OBSERVATORY

# Introduction

- Ultrahigh-energy cosmic rays (UHECRs)
- Extraterrestrial charged particles with ultra-high energies (above  $10^{18}$  eV)
- Acceleration mechanisms and sources are unknown



- Determination of cosmic ray (CR) mass
- Due to scarcity only indirect detection possible
- Estimation relies on quantities statistically encoding mass (number of muons, depth of the shower maximum  $X_{\text{max}}$ )



# Pierre Auger Observatory



- Malargüe, Argentina
- 3000 km<sup>2</sup>

## Surface Detector:

- 1600 stations with a spacing of 1.5 km
- Multiple sub-detectors
- Main detector: water-Cherenkov detectors in a triangular grid




Water-Cherenkov  
Detector




Fluorescence Detector

- Fluorescence Detector:
  - Overlooking the SD
  - Direct  $X_{\text{max}}$  measurement








# Fluorescence detector




- Direct measurement of the longitudinal profile  
→ shower maximum  $X_{\max}$
- 27 telescopes at five sites
  - 6 telescopes at each of four sites
  - 3 telescopes at the High Elevation Auger Telescopes (HEAT) site - lower energies
- Highly sensitive cameras - 15% duty cycle








# Surface detector signal



- Arrival times  
→ shower geometry

- Signal traces from 3 PMTs in each WCD
- Total recorded signal depends on the energy of the primary CR and the distance from the shower core (reconstructed shower axis)
- Shape depends on the different shower components (muonic, electromagnetic)






arXiv:2406.06319v3



# Mass estimation with DNN

- $X_{\max}$  reconstruction from a shower footprint on the SD dataset
  - Vertical events  $\theta < 60^\circ$ ,  $\lg(E/\text{eV}) > 18.5$ , analysis specific selection
  - 48 824 events
- Calibration with the Hybrid dataset
  - Events measured by both SD and FD
  - 1 642 events after high-quality selection





# Investigating the mass composition



- Mass composition of UHECRs connected to the energy evolution of  $\langle X_{\max} \rangle$  and  $\sigma(X_{\max})$
- Using the SD dataset (48 824 events) compared to the full FD dataset (9 822 events)
- $\langle X_{\max} \rangle$  and  $\sigma(X_{\max})$  of SD dataset predictions in agreement with the FD dataset
- Change of  $\langle X_{\max} \rangle$  per decade

of energy described by the *elongation rate*

$$D_{10} = \frac{d\langle X_{\max} \rangle}{d \lg E} = D_{10}^{\text{MC}} \left( 1 - \frac{d\langle \ln A \rangle}{d \ln E} \right)$$

$$\Rightarrow \frac{d\langle \ln A \rangle}{d \ln E} = 1 - \frac{D_{10}}{D_{10}^{\text{MC}}}$$

- Average elongation rate of the SD data agrees with the FD measurements



arXiv:2406.06319v3





# Interpretation of the results

- A simple linear model does not describe the elongation rate well
- Model with 3 breaks with  $4.4\sigma$  significance
  - Non-trivial composition changes
  - Breaks close to the features in the energy spectrum

| Elongation rate |         | Spectrum    |         |
|-----------------|---------|-------------|---------|
| 1st break       | 6.5 EeV | ankle       | 5.1 EeV |
| 2nd break       | 11 EeV  | instep      | 13 EeV  |
| 3rd break       | 30 EeV  | suppression | 48 EeV  |

arXiv:2501.01736v1





# Interpretation of the results



## ■ Converting $X_{\max}$ to $\ln A$ :

$$\langle \ln A \rangle = \frac{\langle X_{\max} \rangle - \langle X_{\max} \rangle_p}{f_E}, \quad V[\ln A] = \frac{\sigma^2(X_{\max}) - \sigma_{\text{sh}}^2(\ln A)}{b\sigma_p^2 - f_E^2},$$


$$f_E = \left( \langle X_{\max} \rangle_{\text{Fe}} - \langle X_{\max} \rangle_p \right) / \ln 56$$

## ■ Light to heavier composition with energy

arXiv-2507.10292v1




arXiv-2507.10292v1






# Summary

- Excellent agreement of DNN  $\langle X_{\max} \rangle$  reconstruction with FD measurements
- Confirming a heavier and purer composition with increasing energy
- A 3-break model fits the elongation data better








PIERRE  
AUGER  
OBSERVATORY

# Back-up



# Mass estimation with DNN

- Triangular to Cartesian grid - 13x13 stations
- Input normalization
- Training with 400 000 simulated detector responses
- CORSIKA with the EPOS-LHC hadronic interaction model (p, He, O, Fe)
- Energy between 1 and 160 EeV, spectral index  $\gamma = -1$ , zenith angles  $\theta < 65^\circ$
- Testing on EPOS-LHC, QGSJetII-04, Sibyll2.3c hadronic interaction models





# Data selection

TABLE I: Basic and analysis-specific selections (separated by a line) for the SD data set.

| Cut                                                     | Events    | $\epsilon$ (%) |
|---------------------------------------------------------|-----------|----------------|
| reconstructed vertical event ( $\theta < 60^\circ$ )    | 5,994,712 | —              |
| is 6T5                                                  | 4,858,291 | 81.1           |
| $\log_{10}(E/\text{eV}) > 18.5$                         | 133,167   | 2.7            |
| hardware status                                         | 129,403   | 97.2           |
| station start slot                                      | 128,308   | 99.2           |
| $2.75 < A/P < 3.45$                                     | 126,033   | 98.2           |
| $\bar{S}_{\text{tot}} > 5$ VEM in surrounding hexagon   | 125,828   | 99.8           |
| $350 \text{ m} < \text{core distance} < 1000 \text{ m}$ | 101,392   | 80.6           |
| fiducial SD cut                                         | 48,824    | 48.1           |

arXiv:2406.06319v3

TABLE II: Selections for hybrid data.





| Cut                    | Events | $\epsilon$ (%) |
|------------------------|--------|----------------|
| number of events       | 25,076 | —              |
| telescope cuts         | 19,733 | 78.7           |
| hardware status        | 16,916 | 85.7           |
| aerosols/clouds        | 9,822  | 58.1           |
| hybrid geometry        | 9,157  | 93.2           |
| fiducial FoV cut       | 3,497  | 38.2           |
| profile cuts           | 3,331  | 95.3           |
| passed SD selection    | 3,086  | 92.6           |
| analysis-specific cuts | 1,642  | 53.2           |

arXiv:2406.06319v3



# Elongation rate models

| Parameter                                   | Const. elong.                                          | 1-break model                                          | 2-break model                                          | 3-break model                                          | Energy spectrum                                        |
|---------------------------------------------|--------------------------------------------------------|--------------------------------------------------------|--------------------------------------------------------|--------------------------------------------------------|--------------------------------------------------------|
|                                             | Val $\pm \sigma_{\text{stat}} \pm \sigma_{\text{sys}}$ | Val $\pm \sigma_{\text{stat}} \pm \sigma_{\text{sys}}$ | Val $\pm \sigma_{\text{stat}} \pm \sigma_{\text{sys}}$ | Val $\pm \sigma_{\text{stat}} \pm \sigma_{\text{sys}}$ | Val $\pm \sigma_{\text{stat}} \pm \sigma_{\text{sys}}$ |
| $b / \text{g cm}^{-2}$                      | $743 \pm 5 \pm 13$                                     | $743 \pm 5 \pm 13$                                     | $750.5 \pm 4 \pm 13$                                   | $750.5 \pm 3 \pm 13$                                   |                                                        |
| $D_0 / \text{g cm}^{-2} \text{decade}^{-1}$ | $24 \pm 1 \pm 4$                                       | $23 \pm 2 \pm 12$                                      | $12 \pm 6 \pm 5$                                       | $12 \pm 5 \pm 6$                                       |                                                        |
| $E_1 / \text{EeV}$                          |                                                        | $35 \pm 12 \pm 16$                                     | $6.5 \pm 0.9 \pm 1$                                    | $6.5 \pm 0.6 \pm 1$                                    | $4.09 \pm 0.1 \pm 0.8$                                 |
| $D_1 / \text{g cm}^{-2} \text{decade}^{-1}$ |                                                        | $39 \pm 14 \pm 12$                                     | $39 \pm 12 \pm 10$                                     | $39 \pm 5 \pm 14$                                      |                                                        |
| $E_2 / \text{EeV}$                          |                                                        |                                                        | $10 \pm 2 \pm 3$                                       | $11 \pm 2 \pm 1$                                       | $14 \pm 1 \pm 2$                                       |
| $D_2 / \text{g cm}^{-2} \text{decade}^{-1}$ |                                                        |                                                        | $22 \pm 3 \pm 8$                                       | $16 \pm 3 \pm 6$                                       |                                                        |
| $E_3 / \text{EeV}$                          |                                                        |                                                        |                                                        | $31 \pm 5 \pm 3$                                       | $47 \pm 3 \pm 6$                                       |
| $D_3 / \text{g cm}^{-2} \text{decade}^{-1}$ |                                                        |                                                        |                                                        | $42 \pm 9 \pm 12$                                      |                                                        |



arXiv:2406.06319v3



$$J(E) = J_0 \left( \frac{E}{10^{16} \text{ eV}} \right)^{-\gamma_0} \prod_{i=0}^4 \left[ 1 + \left( \frac{E}{E_{ij}} \right)^{\frac{1}{\omega_{ij}}} \right]^{(\gamma_i - \gamma_j) \omega_{ij}}, \quad j = i + 1, \quad (1)$$

**Table 1:** Parameters of the best fit of Eq. (1) to the combined spectrum. The first uncertainty is statistical and the second one systematic. The fit has been performed with a set of transition width parameters that well describe the data:  $\omega_{01} = \omega_{12} = 0.25$  and  $\omega_{23} = \omega_{34} = \omega_{45} = 0.05$  [5].

$$J_0 = (8.34 \pm 0.04 \pm 3.40) \times 10^{-11} \text{ km}^{-2} \text{ sr}^{-1} \text{ yr}^{-1} \text{ eV}^{-1}$$

*low energy ankle*  $E_{01} = (2.8 \pm 0.3 \pm 0.4) \times 10^{16} \text{ eV}$

*2<sup>nd</sup> knee*  $E_{12} = (1.58 \pm 0.05 \pm 0.2) \times 10^{17} \text{ eV}$

*ankle*  $E_{23} = (5.0 \pm 0.1 \pm 0.8) \times 10^{18} \text{ eV}$

*instep*  $E_{34} = (1.4 \pm 0.1 \pm 0.2) \times 10^{19} \text{ eV}$

*suppression*  $E_{45} = (4.7 \pm 0.3 \pm 0.6) \times 10^{19} \text{ eV}$

$$\gamma_0 = 3.09 \pm 0.01 \pm 0.10$$

$$\gamma_1 = 2.85 \pm 0.01 \pm 0.05$$

$$\gamma_2 = 3.283 \pm 0.002 \pm 0.10$$

$$\gamma_3 = 2.54 \pm 0.03 \pm 0.05$$

$$\gamma_4 = 3.03 \pm 0.05 \pm 0.10$$

$$\gamma_5 = 5.3 \pm 0.3 \pm 0.1$$

