


Boosting 21-cm Constraints on Dark Matter


Including results from 2 very recent papers, in collaboration with:

- Rouven Essig (YITP, Stony Brook), Daniele Gaggero (INFN, Pisa), Sergio Palomares-Ruiz (IFIC, Valencia), Gregory Suczewski (YITP, Stony Brook), Mauro Valli (INFN, Rome). 2510.14877 — “Astrophysical Uncertainties...”
- Tracy Slatyer (MIT, Cambridge). 2510.26791 — “Boosting the cosmic 21-cm...”


Setting the scene




Setting the scene — Neutral Hydrogen




The 21-cm signal



The 21-cm signal



The 21-cm signal



The 21-cm signal

CMB temperature

Gas kinetic temperature

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_k^{-1}}{1 + x_k + x_\alpha}$$

UV dependent

X-ray dependent

The spin temperature is dictated by its (de)coupling to T_γ and T_K . This is determined by:

1. Absorption and emission of 21cm photons from and to the CMB
2. Collisional coupling between H atoms
3. Wouthuysen-Field Effect (Lyman-alpha photons)

21cm physics (a primer)

Differential brightness temperature

CMB temperature

$$\delta T_b \propto x_{\text{HI}} \left(1 - \frac{T_\gamma}{T_S} \right) \left(\frac{1+z}{10} \right)^{1/2}$$

Fraction of neutral hydrogen

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_\alpha^{-1}}{1 + x_k + x_\alpha}$$

Spin temperature

21cm physics (a primer)

Differential brightness temperature

CMB temperature

$$\delta T_b \propto x_{\text{HI}} \left(1 - \frac{T_\gamma}{T_S} \right) \left(\frac{1+z}{10} \right)^{1/2}$$

Fraction of neutral hydrogen

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_\alpha^{-1}}{1 + x_k + x_\alpha}$$

Spin temperature

- $T_S = T_\gamma \rightarrow \delta T_b = 0$
- $T_S > T_\gamma \rightarrow \delta T_b > 0$
- $T_S < T_\gamma \rightarrow \delta T_b < 0$

21cm physics (a primer)

Differential brightness temperature

CMB temperature

Fraction of neutral hydrogen

Spin temperature

$$\delta T_b \propto x_{\text{HI}} \left(1 - \frac{T_\gamma}{T_S} \right) \left(\frac{1+z}{10} \right)^{1/2}$$


$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_k^{-1}}{1 + x_k + x_\alpha}$$

- $T_S = T_\gamma \rightarrow \delta T_b = 0$
- $T_S > T_\gamma \rightarrow \delta T_b > 0$
- $T_S < T_\gamma \rightarrow \delta T_b < 0$

The signal is dictated by T_γ/T_S !

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_k^{-1}}{1 + x_k + x_\alpha}$$

The 21-cm signal




The 21-cm signal



EDGES




SARAS




HERA



SKA




21cm physics (a primer)



Radiative coupling most efficient (too low densities for collisions) and $T_S \simeq T_\gamma$


21cm physics (a primer)



Radiative coupling most efficient (too low densities for collisions) and $T_S \simeq T_\gamma$

Lyman-alpha photons from first sources couples $T_S \rightarrow T_K$, through the WF effect

21cm physics (a primer)




Radiative coupling most efficient (too low densities for collisions) and $T_S \simeq T_\gamma$

Lyman-alpha photons from first sources couples $T_S \rightarrow T_K$, through the WF effect


X-ray heating of the gas, where WF effect still couples $T_S \rightarrow T_K$. We see emission when $T_K > T_\gamma$

21cm physics (with exotics)



Exotic Injections change two (measurable) things:

1. Extra heating of the gas




Radiative coupling most efficient (too low densities for collisions) and $T_S = T_\gamma$

Lyman-alpha photons from first sources couples $T_S \rightarrow T_K$, through the WF effect

X-ray heating of the gas, where WF effect still couples $T_S \rightarrow T_K$. We see emission when $T_K > T_\gamma$

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_\alpha T_k^{-1} + x_k T_k^{-1}}{1 + x_\alpha + x_k}$$

2. Extra Lyman-alpha photons (new)





Radiative coupling most efficient (too low densities for collisions) and $T_S = T_\gamma$

Lyman-alpha photons from first sources couples $T_S \rightarrow T_K$, through the WF effect

X-ray heating of the gas, where WF effect still couples $T_S \rightarrow T_K$. We see emission when $T_K > T_\gamma$

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_\alpha T_k^{-1} + x_k T_k^{-1}}{1 + x_\alpha + x_k}$$

The signal space from “standard” astrophysics is wide




1709.02122

- redshift of the Ly α coupling
- moment of the heating transition
- midpoint of reionization


Independent constraints on astrophysics

- UVLFs [1809.08995, 2003.04442, 2110.13919, etc.]
- CMB optical depth [1809.08995, 2502.03525, etc.]
- Chandra X-ray and HERA additional heating [2502.03525, 2108.07282, etc.]
- Starburst99 Synthesis models (+ more recent)

Astrophysical parameter	BENCHMARK	LESS-CONSTRAINING	MORE-CONSTRAINING
N_α	9690	3×10^3	4×10^4
$\log_{10} L_X / \dot{M}_* \left[\frac{\text{erg}}{\text{s}} \frac{\text{yr}}{M_\odot} \right]$	40.5	41	39.5
$T_{\text{vir}} [K]$	10^4	10^4	10^5



Accreting PBHs: A typical heating signal





[Image from Jeremy Schnittman - NASA]

$$L = \epsilon (\dot{M}) \dot{M} c^2$$



$$\left. \frac{d^2 E}{dV dt} \right|_{\text{inj}} = L f_{\text{PBH}} \frac{\rho_{\text{DM}}}{M}$$

Details in [2403.18895, 2510.14877]


LESS-CONSTRAINING SCENARIO: $N_\alpha = 3 \times 10^3$, $L_X = 10^{41} \text{ erg s}^{-1} M_\odot^{-1} \text{ yr}$, $T_{\text{vir}} = 10^4 \text{ K}$.



MORE-CONSTRAINING SCENARIO: $N_\alpha = 4 \times 10^4$, $L_X = 10^{39.5} \text{ erg s}^{-1} M_\odot^{-1} \text{ yr}$, $T_{\text{vir}} = 10^5 \text{ K}$.




The astrophysical scenario matters!




Consider Lyman- *α* instead of
heating

Lyman-alpha from particle (e.g. axion) decay



$$\chi \rightarrow \gamma\gamma$$

Lyman-alpha from particle (e.g. axion) decay





$$\chi \rightarrow \gamma\gamma$$

$$T_S^{-1} = \frac{T_\gamma^{-1} + x_k T_k^{-1} + x_\alpha T_k^{-1}}{1 + x_k + x_\alpha}$$

extra Lyman- α (new)


extra heating (old)

Back to the 21-cm signal




2507.10533

Fiducial scenario bound




Fiducial scenario bound





What if astrophysics is “unfavourable”

Scenario	$L_X/\dot{M}_*$	$N_{\text{ion}}^{\text{ACG}}$	$N_{\text{Ly}\alpha}^{\text{ACG}}$	$N_{\text{ion}}^{\text{MCG}}$	$N_{\text{Ly}\alpha}^{\text{MCG}}$
Fiducial	$10^{40.5}$	5×10^3	1.1×10^4	4.4×10^4	4.8×10^3
High L_X	$10^{41.5}$	5×10^3	1.1×10^4	4.4×10^4	4.8×10^3
Low N_α	$10^{40.5}$	1×10^3	2.2×10^3	2×10^4	2.1×10^3
High N_α	$10^{40.5}$	2×10^4	4.4×10^4	8×10^4	8.7×10^3



Fiducial scenario bound






$$m_\chi = 20.4 \text{ eV}$$

	Fiducial	High L_X	High N_α	Low N_α
τ_{95} [s]	1.1×10^{26}	5.2×10^{25}	2.2×10^{26}	1.1×10^{26}



What else can we learn from this?

- Other DM candidates at different scales may emit efficiently in to Lyman-band
- Astrophysics robust probes?
- Dark Ages signal even more robust to these signatures (future lunar detectors)




Summary

- The **uncertainties on the astrophysics**, entering in the null hypothesis, is the **most important systematic** for 21-cm forecasts
- **Lyman-alpha from dark matter can help!**




A1: CMB and 21-cm Signal



$$T_S^{-1} = \frac{T_\gamma^{-1} + x_\alpha T_c^{-1} + x_k T_k^{-1}}{1 + x_\alpha + x_k}$$

$$\delta T_b \simeq 27 \text{ mK } x_{\text{HI}} (1 + \delta) \left(1 - \frac{T_\gamma}{T_S} \right) \left(\frac{1 + z}{10} \right)^{1/2}$$




Pritchard and Loeb (2012)

A3: The EDGES affair


An absorption profile centred at 78 MHz in the sky-averaged spectrum [10.1038/nature25792]

$$\delta T_b \simeq 27 \text{ mK } x_{\text{HI}} (1 + \delta) \left(1 - \frac{T_\gamma}{T_S} \right) \left(\frac{1 + z}{10} \right)^{1/2}$$




EDGES is too low!


Standard scenario with POP II stars



A4: Power spectrum status - HERA and NenuFAR





2210.04912



2507.10533


A5: Systematics related to astrophysics




Virial
Temperature

$$\text{SFR} \propto \exp\left(-\frac{M_{\text{turn}}}{M_h}\right)$$

$$M_{\text{turn}} \propto T_{\text{vir}}^{3/2}$$




X-Ray
Luminosity



Lyman Alpha


[Laura Lopez-Honorez+2016]

A6: BHL



Steady state spherical
H. Bondi (1952)


$$\dot{M}_{\text{BHL}} = 4\pi\lambda \frac{(GM_{\text{PBH}})^2 \rho_b}{(v_{\text{BH}}^2 + c_s^2)^3}$$



Hoyle, F., & Lyttleton, R. A. (1939)

"It seems *likely* that it represents the *order of magnitude* of the accretion rate" - Bondi 1952

A6.1: PR





$$\dot{M}_{\text{PR}} = 4\pi \rho_{\text{in}} \frac{(GM_{\text{PBH}})^2}{(v_{\text{in}}^2 + c_{\text{s,in}}^2)^{3/2}}$$


Fixed by Euler's equations at the ionisation front

Free parameter parametrising temperature in the ionised region

Park and Ricotti 2013: numerical simulations + semi-analytic formula including radiative feedback.




A6.2: PR bound




A7: How to perform a forecast


4. Perform your MCMC, or Fisher Forecast, or compute the likelihood given a fiducial signal, varying only f_{PBH}



1. Implement PBH accretion by modifying
CLASS + ZEUS21



2. Choose a fiducial set of astrophysics parameters for your null hypothesis. We use $(N_\alpha, L_{40}$ and T_{vir}).




3. Define a **Gaussian multivariate** likelihood containing the **null hypothesis**, and sensitivities of SKA at 1080 hours of observation using 21cmSense



$$\log \mathcal{L} = -\frac{1}{2} \sum_{i=1}^n \frac{(x_i - \mu_i^{\text{NULL}})^2}{\sigma_i^2}$$



A8: What else can these Lyman-series photons do?

- Lyman-Werner photons (11.2-13.6 eV) suppress formation of molecular hydrogen
 - ▶ Direct collapse of pristine hydrogen to form SMBHs




2404.03909

A9: Correlations between parameters




A10: X-rays and Lyman-alpha



• X-ray heating of the gas



• Lyman-alpha coupling to the gas



A11: Hera Collaboration 2021+2022



A12: Evolution equations

$$\frac{dX_e(z)}{dz} = \frac{1}{(1+z)H(z)}(R(z) - I(z) - I_X(z))$$

$$\frac{dT_b}{dz} = \frac{1}{1+z} \left[2T_b + \gamma(T_b - T_{CMB}) \right] + K_h$$

$$J_\alpha = J_\alpha^{\text{astro}} + J_\alpha^{\text{DM}}$$