



Unique Properties of Secondary Cosmic Rays Results from the Alpha Magnetic Spectrometer



Jin Zhang

On Behalf of the AMS Collaboration

Shandong University


TeV Particle Astrophysics, Valencia, Spain, Nov, 2025



Introduction

The Nuclei in cosmic rays can be separated into two classes according to their origins: the primary cosmic rays are produced during the lifetime of stars and accelerated in supernovae shocks; The secondary cosmic rays are produced by the collisions of primary nuclei with the interstellar medium.

Measurements of secondary cosmic ray nuclei fluxes are fundamental to understanding the propagation processes of cosmic rays in the Galaxy.



Alpha Magnetic Spectrometer (AMS)


Transition Radiation Detector (TRD)
identify e^+ , e^-




Silicon Tracker
measure Z, P



Ring Imaging Cerenkov (RICH)
measure Z, E



Particle Physics Experiment in Space



Upper TOF measure Z, E



Magnet identify $\pm Z$, P




Anticoincidence Counters (ACC)
reject particles from the side




Lower TOF measure Z, E



AMS Measurement on Cosmic Ray Nuclei Charge



AMS Nuclei Flux Measurements






Measurements require knowledge of detector performance details, the resolution functions, acceptance ... obtained by AMS Monte Carlo Simulations

In AMS 2 to 4 independent analysis are done to compute $N_i, A_i, \varepsilon_i, T_i$ for each flux


Unique of AMS: Measure Nuclear Reaction Cross Section with Cosmic Rays

AMS can collect cosmic particles entering the detector from both the right and the left directions when the detector is flying horizontally. This allows the measurement of reaction cross-sections of various nuclei colliding on the AMS material within the GV to TV rigidity ranges.




Based on the measurements, AMS adjusted the relevant parameters of nuclear interaction models in the simulation data to guarantee a better precision for the spectra measurements.


Secondary Cosmic Rays Selections



Thanks to redundant charge measurements and good charge resolution, AMS is able to identify Li, Be, B and F nuclei with a negligible background (<0.5%) arising from charge misidentification over the whole rigidity range.





Background from Nuclei Interaction in AMS Materials






The background mainly comes from heavier nuclei interaction with AMS material, they are accurately measured (with typical error of few percents). A clean sample is obtained by subtracting the background from measured event numbers.

Example of the measurements of the background arising from the interaction with TRD and UTOF material:




Cosmic Rays Lithium Beryllium and Boron Fluxes







AMS measured the Lithium (Li), Beryllium (Be) and Boron (B) fluxes with unprecedented precision.

Secondary Primary Fluxes Comparison




The secondary Li-Be-B fluxes have the same rigidity dependence above 30 GV, and it is distinctly different with primary cosmic rays.

Secondary to Primary Fluxes Ratios




- Above ~ 200 GV, all three secondary-to-primary flux ratios harden with $\Delta_2 - \Delta_1 = 0.13 \pm 0.025$
- The Δ_2 is not compatible with Δ_1 with C.L more than 5σ show that the spectral hardening at ~ 200 GV is due to propagation effect.

Cosmic Rays Fluorine Flux




AMS measured the Fluorine (F) flux with unprecedented precision and extended the measured rigidity to a much higher range.

Fluorine Spectral Structures



The F/Si flux ratio also show a spectral hardening with $\Delta_2 - \Delta_1 = 0.14 \pm 0.06$ which is compatible with that of B/O flux ratio hardening with 0.13 ± 0.025 .



Above 10 GV, the ratio of $\frac{F/Si}{B/O}$ can be described by a single power law with $\delta = 0.055 \pm 0.006$ ($> 7\sigma$ difference from 0) show that the heavier secondary-to-primary F/Si flux ratio is distinctly different from the light B/O , which indicates that secondary cosmic rays also have two classes.

Summary

- The latest AMS measurements of secondary cosmic ray Li, Be, B and F fluxes are presented
- All fluxes exhibit a hardening above 200 GV, however, the secondary fluxes harden more than the primary fluxes, which favors a propagation origin of the observed hardening
- The $\frac{F/Si}{B/O}$ ratio deviates from a constant by more than 7σ demonstrating that F belongs to different class of secondary cosmic ray with Li-Be-B

