


Quantum Decoherence at Colliders.




Rafael Aoude¹, José Manuel Camacho², Fabio Maltoni³, María Moreno⁴, Leonardo Satrioni⁵, Marcel Vos⁶
Guillermo García⁷

Prospects for $e^+e^- \rightarrow \tau^+\tau^-\gamma$

Entanglement could be seen not only in tops but also in taus at BelleII and a future FCC operating in the Z-pole (Ehataht et al. 2024, M. Fabbrichesi et al. 2024) using hadronic polarimetry.

Can photon decoherence occur in this scenario? Let's explore it!



Offers efficient polarimetry



$$\longrightarrow e^+e^- \rightarrow \tau^+\tau^-\gamma \rightarrow \pi^+\pi^-\nu_\tau\bar{\nu}_\tau\gamma, \quad (\tau^- \rightarrow \pi^-\nu_\tau)$$

Apparently, ISR is not a problem for this type of analysis
(note: suppressed by central region cut at Belle 2, naturally
suppressed at Z-pole)

Prospects for $e^+e^- \rightarrow \tau^+\tau^-\gamma$: Belle II

Experimental setting: Tomography produced for 10M events, Reweight produced for 1M events, non polarized electron beams



Angular cut applied in order to increase the entanglement per bin $\rightarrow \cos(\theta_{\tau^+\tau^-}^{RF}) = \hat{k}_{\tau} * Z \quad Z = (0, 0, 1)$ [in $\tau^+\tau^-$ -rest frame]



No signs of a tauonium singlet entanglement peak induced by hard photon radiation

Prospects for $e^+e^- \rightarrow \tau^+\tau^-\gamma$: Z-pole

Experimental setting: Tomography produced for 10M events, Reweight produced for 1M events, non polarized electron beams





No signs of a tauonium singlet entanglement peak induced by hard photon radiation

Prospects for e^+e^- collisions: pT cut analysis

Now, we will focus on an experimentally-driven analysis, examining how entanglement scales with different pT cuts applied to the FSR (gluon in tops - photon in taus)

Experimental setting: Tomography produced for 1M events, non polarized electron beams





Decoherence exhibits a quasi-linear dependence on the pT cut applied

Prospects for e^+e^- collisions: pT cut analysis

Now, we will focus on an experimentally-driven analysis, examining how entanglement scales with different pT cuts applied to the FSR (gluon in tops - photon in taus)

Experimental setting: Tomography produced for 1M events, non polarized electron beams



Decoherence exhibits a quasi-linear dependence on the pT cut applied

Detection of FSR induced decoherence in e^+e^- collisions

Quantum observables for decoherence

We have defined four observable that match decoherence:

$$\Delta C[\rho] = C[\rho](t\bar{t}) - C[\rho](t\bar{t}g) \longrightarrow \Delta C[\rho] > 0$$

$$R_{C[\rho]} = \frac{C[\rho](t\bar{t}g)}{C[\rho](t\bar{t})} \longrightarrow R_{C[\rho]} < 1$$

Conditions for decoherence

$$\Delta D_n = D_n(t\bar{t}) - D_n(t\bar{t}g) \longrightarrow \Delta D_n > 0$$

$$R_{D_n} = \frac{D_n(t\bar{t}g)}{D_n(t\bar{t})} \longrightarrow R_{D_n} < 1$$

We are going to calculate the significance for this conditions being fulfilled in experimental achievable scenarios

Detection of FSR induced decoherence in e^+e^- collisions

Significance for the detection of these decoherence conditions

$$e^+e^- \rightarrow t\bar{t}$$

Machine	$\int \mathcal{L} dt$ [ab ⁻¹]	Final state	# $t\bar{t}$ events	$t\bar{t}$	$D_n[t\bar{t}]: e^+e^- \rightarrow$ $t\bar{t}g$	$\Delta D_n[t\bar{t}]$ $t\bar{t}g - t\bar{t}$	significance ($\Delta D_n > 0$)	$R_{D_n} = D_n^{t\bar{t}g}/D_n^{t\bar{t}}$	significance ($R_{D_n} < 1$)
linear e^+e^- (LCF) $\sqrt{s} = 500$ GeV	8	ee, $\mu\mu$, $e\mu$	$5 \cdot 10^6 \times 4\%$	-0.421	-0.351 ($p_T > 35\text{GeV}$)	0.071 \pm 0.019	3.67 σ	0.831 \pm 0.045	3.69 σ
	8	$l'l'$ & e,μ + jets	$5 \cdot 10^6 \times 19\%$	idem	idem	idem \pm 0.009	7.35 σ	idem \pm 0.023	7.39 σ
	8	all channels	$5 \cdot 10^6 \times 30\%$	idem	idem	idem \pm 0.008	7.52 σ	idem \pm 0.022	7.55 σ
linear e^+e^- (LCF) $\sqrt{s} = 1$ TeV	8	ee, $\mu\mu$, $e\mu$	$10^6 \times 4\%$	-0.500	-0.343 ($p_T > 175\text{GeV}$)	0.157 \pm 0.036	4.32 σ	0.687 \pm 0.071	4.35 σ
	8	$l'l'$ & e,μ + jets	$10^6 \times 19\%$	idem	idem	idem \pm 0.017	9.39 σ	idem \pm 0.028	9.47 σ
	8	all channels	$10^6 \times 30\%$	idem	idem	idem \pm 0.015	9.98 σ	idem \pm 0.033	10.1 σ

Machine	$\int \mathcal{L} dt$ [ab ⁻¹]	Final state	# $t\bar{t}$ events	$t\bar{t}$	$C[t\bar{t}]: e^+e^- \rightarrow$ $t\bar{t}g$	ΔC $t\bar{t} - t\bar{t}g$	significance ($\Delta C[t\bar{t}] > 0$)	$R = C^{t\bar{t}g}[t\bar{t}]/C^{t\bar{t}}[t\bar{t}]$	significance ($R < 1$)
linear e^+e^- (LCF) $\sqrt{s} = 500$ GeV	8	ee, $\mu\mu$, $e\mu$	$5 \cdot 10^6 \times 4\%$	0.133	0.026 ($p_T > 35\text{GeV}$)	0.105 \pm 0.026	3.81 σ	0.207 \pm 0.194	3.88 σ
	8	$l'l'$ & e,μ + jets	$5 \cdot 10^6 \times 19\%$	idem	idem	idem \pm 0.014	7.27 σ	idem \pm 0.107	7.38 σ
	8	all channels	$5 \cdot 10^6 \times 30\%$	idem	idem	idem \pm 0.013	7.57 σ	idem \pm 0.104	7.59 σ
linear e^+e^- (LCF) $\sqrt{s} = 1$ TeV	8	ee, $\mu\mu$, $e\mu$	$10^6 \times 4\%$	0.248	0.017 ($p_T > 175\text{GeV}$)	0.225 \pm 0.038	5.48 σ	0.068 \pm 0.151	5.64 σ
	8	$l'l'$ & e,μ + jets	$10^6 \times 19\%$	idem	idem	idem \pm 0.0225	10.3 σ	idem \pm 0.083	11.02 σ
	8	all channels	$10^6 \times 30\%$	idem	idem	idem \pm 0.021	10.48 σ	idem \pm 0.078	11.6 σ

Detection of FSR induced decoherence in e^+e^- collisions

Significance for the detection of these decoherence conditions

$$\underline{e^+e^- \rightarrow \tau^+\tau^- (\tau^- \rightarrow \pi^-\nu_\tau)}$$

Machine	\sqrt{s} [GeV]	$\int \mathcal{L} dt$ [ab ⁻¹]	# $t\bar{t}$ events	$D_n[\tau^+\tau^-]: e^+e^- \rightarrow$ $\tau^+\tau^-$ $\tau^+\tau^-\gamma$		$\Delta D_n[\tau^+\tau^-]$ $\tau^+\tau^-\gamma - \tau^+\tau^-$	significance ($\Delta D_n > 0$)	$R_{D_n} = D_n^{\tau^+\tau^-\gamma} / D_n^{\tau^+\tau^-}$	significance ($R_{D_n} < 1$)
Belle II 2025	10.579	0.5	5×10^8	-0.818	-0.488 ($p_T > 2\text{GeV}$)	0.331 ± 0.011	28.8 σ	0.596 ± 0.014	28.9 σ
Belle II Tarjet	10.579	50	5×10^{10}	idem	idem	idem ± 0.002	288 σ	idem ± 0.001	289 σ
Z-pole (GigaZ) GeV	91.2	0.1	10^8	-0.911	-0.5846 ($p_T > 20\text{GeV}$)	0.327 ± 0.018	17.5 σ	0.641	17.54 σ
Z-pole (TeraZ)	91.2	300	2×10^{11}	idem	idem	idem ± 0.0006	958 σ	idem ± 0.0003	960 σ

Machine	\sqrt{s} [GeV]	$\int \mathcal{L} dt$ [ab ⁻¹]	# $\tau^+\tau^-$ events	$C[\tau^+\tau^-]: e^+e^- \rightarrow$ $\tau^+\tau^-$ $\tau^+\tau^-\gamma$		$\Delta C[\tau^+\tau^-]$ $\tau^+\tau^- - \tau^+\tau^-\gamma$	significance ($\Delta C[\tau^+\tau^-] > 0$)	$R = C^{\tau^+\tau^-\gamma} / C^{\tau^+\tau^-}$	significance ($R < 1$)
Belle II 2025	10.579	0.5	5×10^8	0.726	0.234 ($p_T > 2\text{GeV}$)	0.492 ± 0.017	27.2 σ	$0.323 \pm$	27.9 σ
Belle II Tarjet	10.579	50	5×10^{10}	idem	idem	idem ± 0.002	277 σ	idem ± 0.002	279 σ
Z-pole (GigaZ) GeV	91.2	0.1	10^8	0.858	0.381 ($p_T > 20\text{GeV}$)	0.477 ± 0.029	16.32 σ	0.444 ± 0.034	16.40 σ
Z-pole (TeraZ)	91.2	300	2×10^{11}	idem	idem	idem ± 0.0005	895 σ	idem ± 0.001	864 σ

Thanks for your attention!!

